

# Non-linear problems in $n$ variable

Lectures for PHD course on  
Non-linear equations and numerical optimization

Enrico Bertolazzi  
DIMS – Università di Trento  
March 2005



## Outline

- 1 The Newton Raphson
- 2 The Broyden method
- 3 The dumped Broyden method



## The problem to solve

### Problem

Given  $\mathbf{F} : D \subseteq \mathbb{R}^n \mapsto \mathbb{R}^n$   
Find  $\mathbf{x}_* \in D$  for which  $\mathbf{F}(\mathbf{x}_*) = \mathbf{0}$ .

### Example

Let

$$\mathbf{F}(\mathbf{x}) = \begin{pmatrix} x_1^2 + x_2^3 + 7 \\ x_1 + x_2 + 1 \end{pmatrix}$$

which has  $\mathbf{F}(\mathbf{x}_*) = \mathbf{0}$  for  $\mathbf{x}_* = (1, -2)^T$ .



## The Newton Raphson

## Outline

- 1 The Newton Raphson
- 2 The Broyden method
- 3 The dumped Broyden method



## The Newton procedure

(1/3)

- Consider the following map

$$\mathbf{F}(\mathbf{x}) = \begin{pmatrix} x_1^2 + x_2^3 + 7 \\ x_1 + x_2 + 1 \end{pmatrix}$$

we know an approximation of a root  $\mathbf{x}_0 \approx (1.1, -1.9)^T$ .

- Setting  $\mathbf{x}_1 = \mathbf{x}_0 + \mathbf{p}$  we obtain <sup>1</sup>

$$\mathbf{F}(\mathbf{x}_0 + \mathbf{p}) = \begin{pmatrix} 1.351 \\ 0.2 \end{pmatrix} + \begin{pmatrix} 2.2 & 10.83 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} p_1 \\ p_2 \end{pmatrix} + \mathcal{O}(\|\mathbf{p}\|^2)$$

if  $\mathbf{x}_0$  is a good approximation of a root of  $\mathbf{F}(\mathbf{x})$  then  $\mathcal{O}(\|\mathbf{p}\|^2)$  is a small vector.

<sup>1</sup>Here  $\mathcal{O}(\mathbf{x})$  means  $(\mathcal{O}(x_1), \dots, \mathcal{O}(x_n))^T$



## The Newton procedure

(2/3)

- Neglecting  $\mathcal{O}(\|\mathbf{p}\|^2)$  and solving

$$\begin{pmatrix} 1.351 \\ 0.2 \end{pmatrix} + \begin{pmatrix} 2.2 & 10.83 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} p_1 \\ p_2 \end{pmatrix} = \mathbf{0}$$

we obtain  $\mathbf{p} = (-0.094438, -0.105562)^T$ .

- Now we set

$$\mathbf{x}_1 = \mathbf{x}_0 + \mathbf{p} = \begin{pmatrix} 1.005562 \\ -2.0055612 \end{pmatrix}$$



## The Newton procedure

(3/3)

- Considering

$$\mathbf{F}(\mathbf{x}_1 + \mathbf{q}) = \begin{pmatrix} -0.05576 \\ 8 \cdot 10^{-7} \end{pmatrix} + \begin{pmatrix} 2.0111 & 12.0668 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} q_1 \\ q_2 \end{pmatrix} + \mathcal{O}(\|\mathbf{q}\|^2)$$

- Neglecting  $\mathcal{O}(\|\mathbf{q}\|^2)$  and solving

$$\begin{pmatrix} -0.05576 \\ 8 \cdot 10^{-7} \end{pmatrix} + \begin{pmatrix} 2.0111 & 12.0668 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} q_1 \\ q_2 \end{pmatrix} = \mathbf{0}$$

we obtain  $\mathbf{q} = (-0.0055466, 0.0055458)^T$ .

- Now we set  $\mathbf{x}_2 = \mathbf{x}_1 + \mathbf{q} = (1.000015, -2.000015)^T$



## The Newton procedure: a modern point of view

(1/2)

The previous procedure can be resumed as follows:

- Consider the following function  $\mathbf{F}(\mathbf{x})$ . We know an approximation of a root  $\mathbf{x}_0$ .
- Expand by Taylor series

$$\mathbf{F}(\mathbf{x}) = \mathbf{F}(\mathbf{x}_0) + \nabla \mathbf{F}(\mathbf{x}_0)(\mathbf{x} - \mathbf{x}_0) + \mathcal{O}(\|\mathbf{x} - \mathbf{x}_0\|^2)$$

- Drop the term  $\mathcal{O}(\|\mathbf{x} - \mathbf{x}_0\|^2)$  and solve

$$\mathbf{0} = \mathbf{F}(\mathbf{x}_0) + \nabla \mathbf{F}(\mathbf{x}_0)(\mathbf{x} - \mathbf{x}_0)$$

Call  $\mathbf{x}_1$  this solution.

- Repeat 1 – 3 with  $\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots$



## The Newton procedure: a modern point of view (2/2)

## Algorithm (Newton iterative scheme)

Let  $x_0$  assigned, then for  $k = 0, 1, 2, \dots$

- 1 Solve for  $p_k$ :

$$\nabla \mathbf{F}(x_k) p_k + \mathbf{F}(x_k) = 0$$

- 2 Update

$$x_{k+1} = x_k + p_k$$



## Proof.

From basic Calculus:

$$\mathbf{F}(y) - \mathbf{F}(x) = \int_0^1 \nabla \mathbf{F}(x + t(y-x))(y-x) dt$$

subtracting on both side  $\nabla \mathbf{F}(x)(y-x)$  we have

$$\begin{aligned} \mathbf{F}(y) - \mathbf{F}(x) - \nabla \mathbf{F}(x)(y-x) &= \\ \int_0^1 [\nabla \mathbf{F}(x + t(y-x)) - \nabla \mathbf{F}(x)](y-x) dt & \end{aligned}$$

and taking the norm

$$\|\mathbf{F}(y) - \mathbf{F}(x) - \nabla \mathbf{F}(x)(y-x)\| \leq \int_0^1 \gamma t \|y-x\|^2 dt$$



## Standard Assumptions

In the study of convergence of numerical scheme, some standard regularity assumption are assumed for the function  $\mathbf{F}(x)$ .

## Assumption (Standard Assumptions)

The function  $\mathbf{F} : D \subset \mathbb{R}^n \mapsto \mathbb{R}^n$  is continuous, differentiable with Lipschitz derivative  $\nabla \mathbf{F}(x)$ . i.e.

$$\|\nabla \mathbf{F}(x) - \nabla \mathbf{F}(y)\| \leq \gamma \|x - y\| \quad \forall x, y \in D \subset \mathbb{R}^n$$

## Lemma (Taylor like expansion)

Let  $\mathbf{F}(x)$  satisfy the standard assumptions, then

$$\|\mathbf{F}(y) - \mathbf{F}(x) - \nabla \mathbf{F}(x)(y-x)\| \leq \frac{\gamma}{2} \|x-y\|^2 \quad \forall x, y \in D \subset \mathbb{R}^n$$



## Lemma (Jacobian norm control)

Let  $\mathbf{F}(x)$  satisfying standard assumptions, and  $\nabla \mathbf{F}(x_*)$  non singular. Then there exists  $\delta > 0$  such that for all  $\|x - x_*\| \leq \delta$  we have

$$2^{-1} \|\nabla \mathbf{F}(x)\| \leq \|\nabla \mathbf{F}(x_*)\| \leq 2 \|\nabla \mathbf{F}(x)\|$$

and

$$2^{-1} \|\nabla \mathbf{F}(x)^{-1}\| \leq \|\nabla \mathbf{F}(x_*)^{-1}\| \leq 2 \|\nabla \mathbf{F}(x)^{-1}\|$$



Proof.

(1/3).

From standard assumptions choosing  $\gamma\delta \leq 2^{-1} \|\nabla\mathbf{F}(\mathbf{x}_*)\|$

$$\begin{aligned}\|\nabla\mathbf{F}(\mathbf{x})\| &\leq \|\nabla\mathbf{F}(\mathbf{x}) - \nabla\mathbf{F}(\mathbf{x}_*)\| + \|\nabla\mathbf{F}(\mathbf{x}_*)\| \\ &\leq \gamma \|\mathbf{x} - \mathbf{x}_*\| + \|\nabla\mathbf{F}(\mathbf{x}_*)\| \\ &\leq (3/2) \|\nabla\mathbf{F}(\mathbf{x}_*)\| \leq 2 \|\nabla\mathbf{F}(\mathbf{x}_*)\|\end{aligned}$$

again choosing  $\gamma\delta \leq 2^{-1} \|\nabla\mathbf{F}(\mathbf{x}_*)\|$

$$\begin{aligned}\|\nabla\mathbf{F}(\mathbf{x}_*)\| &\leq \|\nabla\mathbf{F}(\mathbf{x}_*) - \nabla\mathbf{F}(\mathbf{x})\| + \|\nabla\mathbf{F}(\mathbf{x})\| \\ &\leq \gamma \|\mathbf{x} - \mathbf{x}_*\| + \|\nabla\mathbf{F}(\mathbf{x})\| \\ &\leq 2^{-1} \|\nabla\mathbf{F}(\mathbf{x}_*)\| + \|\nabla\mathbf{F}(\mathbf{x})\|\end{aligned}$$

so that  $2^{-1} \|\nabla\mathbf{F}(\mathbf{x}_*)\| \leq \|\nabla\mathbf{F}(\mathbf{x})\|$ .



Proof.

(3/3).

Using last inequality again

$$\begin{aligned}\|\nabla\mathbf{F}(\mathbf{x})^{-1}\| &\leq \|\nabla\mathbf{F}(\mathbf{x})^{-1} - \nabla\mathbf{F}(\mathbf{x}_*)^{-1}\| + \|\nabla\mathbf{F}(\mathbf{x}_*)^{-1}\| \\ &\leq 2^{-1} \|\nabla\mathbf{F}(\mathbf{x}_*)^{-1}\| + \|\nabla\mathbf{F}(\mathbf{x}_*)^{-1}\|\end{aligned}$$

so that

$$2^{-1} \|\nabla\mathbf{F}(\mathbf{x})^{-1}\| \leq \|\nabla\mathbf{F}(\mathbf{x}_*)^{-1}\|$$

choosing  $\delta$  such that for all  $\|\mathbf{x} - \mathbf{x}_*\| \leq \delta$  we have  $\nabla\mathbf{F}(\mathbf{x})$  non singular and  $\gamma\delta \leq 2^{-1} \|\nabla\mathbf{F}(\mathbf{x}_*)\|$  and  $\gamma\delta \|\nabla\mathbf{F}(\mathbf{x}_*)^{-1}\| \leq 2^{-1}$  then the inequality of the lemma are true.  $\square$



Proof.

(2/3).

From the continuity of the determinant there exists a neighbor with  $\nabla\mathbf{F}(\mathbf{x})$  non singular for all  $\|\mathbf{x} - \mathbf{x}_*\| \leq \delta$ .

$$\begin{aligned}\|\nabla\mathbf{F}(\mathbf{x})^{-1} - \nabla\mathbf{F}(\mathbf{x}_*)^{-1}\| \\ &\leq \|\nabla\mathbf{F}(\mathbf{x})^{-1}\| \|\nabla\mathbf{F}(\mathbf{x}_*) - \nabla\mathbf{F}(\mathbf{x})\| \|\nabla\mathbf{F}(\mathbf{x}_*)^{-1}\| \\ &\leq \gamma \|\mathbf{x} - \mathbf{x}_*\| \|\nabla\mathbf{F}(\mathbf{x})^{-1}\| \|\nabla\mathbf{F}(\mathbf{x}_*)^{-1}\|\end{aligned}$$

and choosing  $\delta$  such that  $\gamma\delta \|\nabla\mathbf{F}(\mathbf{x}_*)^{-1}\| \leq 2^{-1}$  we have

$$\|\nabla\mathbf{F}(\mathbf{x})^{-1} - \nabla\mathbf{F}(\mathbf{x}_*)^{-1}\| \leq 2^{-1} \|\nabla\mathbf{F}(\mathbf{x}_*)^{-1}\|$$

and using this last inequality

$$\begin{aligned}\|\nabla\mathbf{F}(\mathbf{x}_*)^{-1}\| &\leq \|\nabla\mathbf{F}(\mathbf{x}_*)^{-1} - \nabla\mathbf{F}(\mathbf{x})^{-1}\| + \|\nabla\mathbf{F}(\mathbf{x})^{-1}\| \\ &\leq (3/2) \|\nabla\mathbf{F}(\mathbf{x}_*)^{-1}\| \leq 2 \|\nabla\mathbf{F}(\mathbf{x}_*)^{-1}\|\end{aligned}$$



### Theorem (Local Convergence of Newton method)

Let  $\mathbf{F}(\mathbf{x})$  satisfying standard assumptions, and  $\mathbf{x}_*$  a simple root (i.e.  $\nabla\mathbf{F}(\mathbf{x}_*)$  non singular). Then, if  $\|\mathbf{x}_0 - \mathbf{x}_*\| \leq \delta$  with  $C\delta \leq 1$  where

$$C = \gamma \|\nabla\mathbf{F}(\mathbf{x}_*)^{-1}\|$$

then, the sequence generated by Newton method satisfies:

- 1.  $\|\mathbf{x}_k - \mathbf{x}_*\| \leq \delta$  for  $k = 0, 1, 2, 3, \dots$
- 2.  $\|\mathbf{x}_{k+1} - \mathbf{x}_*\| \leq C \|\mathbf{x}_k - \mathbf{x}_*\|^2$  for  $k = 0, 1, 2, 3, \dots$
- 3.  $\lim_{k \rightarrow \infty} \mathbf{x}_k = \mathbf{x}_*$ .

- The point 2 of the theorem is the second  $q$ -order of convergence of Newton method.



## Proof.

Consider a Newton step with  $\|x_k - x_*\| \leq \delta$  and

$$\begin{aligned}x_{k+1} - x_* &= x_k - x_* - \nabla F(x_k)^{-1} [F(x_k) - F(x_*)] \\ &= \nabla F(x_k)^{-1} [\nabla F(x_k)(x_k - x_*) - F(x_k) + F(x_*)]\end{aligned}$$

taking the norm and using Taylor like lemma

$$\|x_{k+1} - x_*\| \leq 2^{-1}\gamma \|x_k - x_*\|^2 \|\nabla F(x_k)^{-1}\|$$

from **Jacobian norm control** lemma there exist a  $\delta$  such that  $2 \|\nabla F(x_k)^{-1}\| \geq \|\nabla F(x_*)^{-1}\|$  for all  $\|x_k - x_*\| \leq \delta$ . Reducing eventually  $\delta$  such that  $\gamma \delta \|\nabla F(x_*)^{-1}\| \leq 1$  we have

$$\|x_{k+1} - x_*\| \leq \gamma \|\nabla F(x_*)^{-1}\| \delta \|x_k - x_*\|^2 \leq \|x_k - x_*\|,$$

So that by induction we prove point 1. Point 2 and 3 follows trivially.  $\square$

- We can apply for example the gradient method to the merit function  $f(x)$ . This produce a slow method.
- Instead, we can use the Newton method to produce a search direction. The resulting method is the following
  - 1 Compute the search direction by solving  $\nabla F(x_k)d_k + F(x_k) = \mathbf{0}$ ;
  - 2 Find an approximate solution of the problem  $\alpha_k = \arg \min_{\alpha \geq 0} \|\mathbf{F}(x_k + \alpha d_k)\|^2$ ;
  - 3 Update the solution  $x_{k+1} = x_k + \alpha_k d_k$ .
- The previous algorithm **work** if the direction  $d_k$  is a **descent direction**.

- The problem of Newton method is that it converge normally only when  $x_0$  is near  $x_*$  a root of the nonlinear system.
- A way to make a more robust non linear solver is to use the techniques developed for minimization to make a **globally convergent** nonlinear solver.
- In particular if we consider the **merit function**

$$f(x) = \frac{1}{2} \|\mathbf{F}(x)\|^2$$

we have that  $f(x) \geq 0$  and if  $x_*$  is such that  $f(x_*) = 0$  than we have that

- 1  $x_*$  is a global minimum of  $f(x)$ ;
- 2  $\mathbf{F}(x_*) = \mathbf{0}$ , i.e. is a solution of the nonlinear system  $\mathbf{F}(x)$ .
- So that finding a global minimum of the **merit function**  $f(x)$  is the same of finding a solution of the nonlinear system  $\mathbf{F}(x)$ .

Is  $d_k$  a descent direction?

(1/2)

Consider the gradient of  $f(x) = (1/2) \|\mathbf{F}(x)\|^2$ :

$$\begin{aligned}\frac{\partial}{\partial x_k} f(x) &= \frac{1}{2} \frac{\partial}{\partial x_k} \|\mathbf{F}(x)\|^2 = \frac{1}{2} \frac{\partial}{\partial x_k} \sum_{i=1}^n F_i(x)^2 \\ &= \sum_{i=1}^n \frac{\partial F_i(x)}{\partial x_k} F_i(x)\end{aligned}$$

this can be written as

$$\nabla f(x) = \mathbf{F}(x)^T \nabla \mathbf{F}(x)$$

Is  $d_k$  a descent direction?

(2/2)

Now we check  $\nabla f(x_k)d_k$ :

$$\begin{aligned}\nabla f(x_k)d_k &= \mathbf{F}(x_k)^T \nabla \mathbf{F}(x_k)d_k \\ &= -\mathbf{F}(x_k)^T \nabla \mathbf{F}(x_k) \nabla \mathbf{F}(x_k)^{-1} \mathbf{F}(x_k) \\ &= -\mathbf{F}(x_k)^T \mathbf{F}(x_k) \\ &= -\|\mathbf{F}(x_k)\|^2 < 0\end{aligned}$$

so that **Newton direction** is a descent direction.



## Algorithm (The globalized Newton method)

```

k ← 0; x assigned;
f ← F(x);
while ||f_k|| > ε do
  — Evaluate search direction
  Solve ∇F(x)d = F(x);
  — Evaluate dumping factor λ
  Approximate λ = arg min_{α>0} ||F(x - αd_k)||^2 by line-search;
  — perform step
  x ← x - λd;
  f ← F(x);
  k ← k + 1;
end while
  
```

Is the angle from  $d_k$  and  $\nabla f(x_k)$  bounded from  $\pi/2$ ? (2/2)

Let  $\theta_k$  the angle form  $\nabla f(x_k)$  and  $d_k$ , then we have

$$\begin{aligned}\cos \theta_k &= -\frac{\nabla f(x_k)d_k}{\|\mathbf{F}(x_k)\| \|\nabla \mathbf{F}(x_k)^{-1} \mathbf{F}(x_k)\|} \\ &= \frac{\|\mathbf{F}(x_k)\|}{\|\nabla \mathbf{F}(x_k)^{-1} \mathbf{F}(x_k)\|} \\ &\geq \frac{\|\mathbf{F}(x_k)\|}{\|\nabla \mathbf{F}(x_k)^{-1}\| \|\mathbf{F}(x_k)\|} \\ &\geq \|\nabla \mathbf{F}(x_k)^{-1}\|^{-1}\end{aligned}$$

so that, if for example  $\|\nabla \mathbf{F}(x)^{-1}\|$  is bounded from below then the angle  $\theta_k$  is strictly less then  $\pi/2$  radians. By the Zoutendijk theorem then the **globalized Newton scheme** is globally convergent.



## Outline

- 1 The Newton Raphson
- 2 The Broden method
- 3 The dumped Broden method



## The Broyden method

(1/5)

- Newton method is a **fast** ( $q$ -order 2) numerical scheme to approximate the root of a function  $\mathbf{F}(\mathbf{x})$  but needs the knowledge of the Jacobian  $\nabla\mathbf{F}(\mathbf{x})$ .
- Sometimes Jacobian is not available or too expensive to compute, in this case a numerical procedure to approximate the root which does not use derivative is mandatory.
- The Newton scheme find successively the root of the affine approximation

$$L_k(\mathbf{x}) \doteq \nabla\mathbf{F}(\mathbf{x}_k)(\mathbf{x} - \mathbf{x}_k) + \mathbf{F}(\mathbf{x}_k) = \mathbf{0}$$

- Substituting the Jacobian in the affine approximation by  $\mathbf{A}_k$

$$M_k(\mathbf{x}) \doteq \mathbf{A}_k(\mathbf{x} - \mathbf{x}_k) + \mathbf{F}(\mathbf{x}_k) = \mathbf{0}$$

and solving successively this **affine model** produces the family of different methods:



## The Broyden method

(2/5)

## Algorithm (Generic Secant iterative scheme)

Let  $\mathbf{x}_0$  and  $\mathbf{A}_0$  assigned, then for  $k = 0, 1, 2, \dots$

- Solve for  $\mathbf{p}_k$ :

$$M_k(\mathbf{p}_k + \mathbf{x}_k) = \mathbf{A}_k\mathbf{p}_k + \mathbf{F}(\mathbf{x}_k) = \mathbf{0}$$

- Update the root approximation

$$\mathbf{x}_{k+1} = \mathbf{x}_k + \mathbf{p}_k$$

- Update the affine model and produce  $\mathbf{A}_{k+1}$ .



## The Broyden method

(3/5)

- The way an update of  $M_k \rightarrow M_{k+1}$  determine the algorithm.
- A simple update is the forcing of a number of the **secant** relation:

$$M_{k+1}(\mathbf{x}_{k+1-\ell}) = \mathbf{F}(\mathbf{x}_{k+1-\ell}), \quad \ell = 1, 2, \dots, m$$

notice that  $M_{k+1}(\mathbf{x}_{k+1}) = \mathbf{F}(\mathbf{x}_{k+1})$  for all  $\mathbf{A}_{k+1}$ .

- If  $\mathbf{A}_{k+1} \in \mathbb{R}^{n \times n}$  and  $m = n$  and  $\mathbf{d}_\ell = \mathbf{x}_{k+1-\ell} - \mathbf{x}_{k+1}$  are linearly independent then we have enough linear relation to determine  $\mathbf{A}_{k+1}$ .
- Unfortunately vectors  $\mathbf{d}_\ell$  tends to become linearly dependent so that this approach is very ill conditioned.
- A more feasible approach uses less **secant** relation and others conditions to determine  $M_{k+1}$ .



## The Broyden method

(4/5)

- The way an update of  $M_k \rightarrow M_{k+1}$  in Broyden scheme is the following:
  - $M_{k+1}(\mathbf{x}_k) = \mathbf{F}(\mathbf{x}_k)$ ;
  - $M_{k+1}(\mathbf{x}) - M_k(\mathbf{x})$  is small in some sense;
- The first condition imply

$$\mathbf{A}_{k+1}(\mathbf{x}_k - \mathbf{x}_{k+1}) + \mathbf{F}(\mathbf{x}_{k+1}) = \mathbf{F}(\mathbf{x}_k)$$

which set  $n$  linear equation that do not determine the  $n^2$  coefficients of  $\mathbf{A}_{k+1}$ .

- The second condition become

$$M_{k+1}(\mathbf{x}) - M_k(\mathbf{x}) = (\mathbf{A}_{k+1} - \mathbf{A}_k)(\mathbf{x} - \mathbf{x}_k)$$

$$\|M_{k+1}(\mathbf{x}) - M_k(\mathbf{x})\| \leq \|\mathbf{A}_{k+1} - \mathbf{A}_k\| \|\mathbf{x} - \mathbf{x}_k\|$$

where  $\|\cdot\|$  is some norm. The term  $\|\mathbf{x} - \mathbf{x}_k\|$  is not controllable, so a condition should be  $\|\mathbf{A}_{k+1} - \mathbf{A}_k\|$  is minimum.



## The Broyden method

(5/5)

## Defining

$$\mathbf{y}_k = \mathbf{F}(\mathbf{x}_{k+1}) - \mathbf{F}(\mathbf{x}_k), \quad \mathbf{s}_k = \mathbf{x}_{k+1} - \mathbf{x}_k$$

the Broyden scheme find the update  $\mathbf{A}_{k+1}$  which satisfy:

- $\mathbf{A}_{k+1}\mathbf{s}_k = \mathbf{y}_k$ ;
  - $\|\mathbf{A}_{k+1} - \mathbf{A}_k\| \leq \|\mathbf{B} - \mathbf{A}_k\|$  for all  $\mathbf{B}$  such that  $\mathbf{B}\mathbf{s}_k = \mathbf{y}_k$ .
- If we choose for the norm  $\|\cdot\|$  the Frobenius norm  $\|\cdot\|_F$

$$\|\mathbf{A}\|_F = \left( \sum_{i,j=1}^n A_{ij}^2 \right)^{1/2}$$

then the problem admits a unique solution.



## The Frobenius matrix norm

(2/4)

The first two point of the Frobenius norm  $\|\cdot\|_F$  are trivial, to prove point 3 and 4 we need two classical inequality:

## Cauchy-Schwartz inequality

$$\sum_{i=1}^n a_i b_i \leq \left( \sum_{i=1}^n a_i^2 \right)^{1/2} \left( \sum_{i=1}^n b_i^2 \right)^{1/2}$$

The inequality is strict unless  $a_i = \lambda b_i$  for  $i = 1, 2, \dots, n$ .

## Triangular inequality

$$\left( \sum_{i=1}^n (a_i + b_i)^2 \right)^{1/2} \leq \left( \sum_{i=1}^n a_i^2 \right)^{1/2} + \left( \sum_{i=1}^n b_i^2 \right)^{1/2}$$

The inequality is strict unless  $a_i = \lambda b_i$  for  $i = 1, 2, \dots, n$ .



## The Frobenius matrix norm

(1/4)

The Frobenius norm  $\|\cdot\|_F$

$$\|\mathbf{A}\|_F = \left( \sum_{i,j=1}^n A_{ij}^2 \right)^{1/2}$$

is a matrix norm, i.e. it satisfy:

- $\|\mathbf{A}\|_F \geq 0$  and  $\|\mathbf{A}\|_F = 0 \iff \mathbf{A} = \mathbf{0}$ ;
- $\|\lambda\mathbf{A}\|_F = |\lambda| \|\mathbf{A}\|_F$ ;
- $\|\mathbf{A} + \mathbf{B}\|_F \leq \|\mathbf{A}\|_F + \|\mathbf{B}\|_F$ ;
- $\|\mathbf{A}\mathbf{B}\|_F \leq \|\mathbf{A}\|_F \|\mathbf{B}\|_F$ ;

The Frobenius norm is the **length** of the vector  $\mathbf{A}$  if we consider  $\mathbf{A}$  as a vector in  $\mathbb{R}^{n^2}$ .



## The Frobenius matrix norm

(3/4)

Proof of  $\|\mathbf{A} + \mathbf{B}\|_F \leq \|\mathbf{A}\|_F + \|\mathbf{B}\|_F$ .  
By using triangular inequality

$$\begin{aligned} \|\mathbf{A} + \mathbf{B}\|_F &= \left( \sum_{i,j=1}^n (A_{ij} + B_{ij})^2 \right)^{1/2} \\ &\leq \left( \sum_{i,j=1}^n A_{ij}^2 \right)^{1/2} + \left( \sum_{i,j=1}^n B_{ij}^2 \right)^{1/2} \\ &= \|\mathbf{A}\|_F + \|\mathbf{B}\|_F. \end{aligned}$$





## The Frobenius matrix norm

(4/4)

Proof of  $\|AB\|_F \leq \|A\|_F \|B\|_F$ .

By using Cauchy-Schwartz inequality with

$$\begin{aligned} \|AB\|_F &= \left( \sum_{i,j=1}^n \left( \sum_{k=1}^n A_{ik} B_{kj} \right)^2 \right)^{1/2} \\ &\leq \left( \sum_{i,j=1}^n \left( \sum_{k=1}^n A_{ik}^2 \right) \left( \sum_{k'=1}^n B_{k'j}^2 \right) \right)^{1/2} \\ &= \left( \left( \sum_{i=1}^n \sum_{k=1}^n A_{ik}^2 \right) \left( \sum_{j=1}^n \sum_{k'=1}^n B_{k'j}^2 \right) \right)^{1/2} \\ &= \|A\|_F \|B\|_F. \end{aligned}$$



With the Frobenius matrix norm it is possible to solve the following problem

## Lemma

Let  $A \in \mathbb{R}^{n \times n}$  and  $s, y \in \mathbb{R}^n$  with  $s \neq 0$ . Consider the set

$$B = \{B \in \mathbb{R}^{n \times n} \mid Bs = y\}$$

then there exists a **unique** matrix  $B \in \mathcal{B}$  such that

$$\|A - B\|_F \leq \|A - C\|_F \quad \text{for all } C \in \mathcal{B}$$

moreover  $B$  has the following form

$$B = A + \frac{(y - As)s^T}{s^T s}$$

i.e.  $B$  is a rank one perturbation of the matrix  $A$ .

## Proof.

(1/4).

First of all notice that  $\mathcal{B}$  is not empty, in fact

$$\frac{1}{s^T s} y s^T \in \mathcal{B} \quad \left[ \frac{1}{s^T s} y s^T \right] s = y$$

So that the problem is not empty. Next we reformulate the problem as a constrained minimum problem:

$$\arg \min_{B \in \mathbb{R}^{n \times n}} \frac{1}{2} \sum_{i,j=1}^n (A_{ij} - B_{ij})^2 \quad \text{subject to } Bs = y.$$

The solution is a stationary point of the Lagrangian:

$$g(B, \lambda) = \frac{1}{2} \sum_{i,j=1}^n (A_{ij} - B_{ij})^2 + \sum_{i=1}^n \lambda_i \left( \sum_{j=1}^n B_{ij} s_j - y_i \right)$$



## Proof.

(2/4).

taking the gradient we have

$$\frac{\partial}{\partial B_{ij}} g(B, \lambda) = A_{ij} - B_{ij} + \lambda_i s_j = 0$$

$$\frac{\partial}{\partial \lambda_i} g(B, \lambda) = \sum_{j=1}^n B_{ij} s_j - y_i = 0$$

The previous equality can be written in matrix form

$$B = A + \lambda s^T \quad Bs = y$$

so that we can solve for  $\lambda$ 

$$Bs = As + \lambda s^T s = y \quad \lambda = \frac{y - As}{s^T s}$$

next we prove that  $B$  is the **unique minimum**.

**Proof.** (3/4).

The matrix  $B$  is a minimum, in fact

$$\|B - A\|_F = \left\| A + \frac{(y - As)s^T}{s^T s} - A \right\|_F = \left\| \frac{(y - As)s^T}{s^T s} \right\|_F$$

for all  $C \in \mathcal{B}$  we have  $Cs = y$  so that

$$\begin{aligned} \|B - A\|_F &= \left\| \frac{(Cs - As)s^T}{s^T s} \right\|_F = \left\| (C - A) \frac{ss^T}{s^T s} \right\|_F \\ &\leq \|C - A\|_F \left\| \frac{ss^T}{s^T s} \right\|_F = \|C - A\|_F \end{aligned}$$

because in general

$$\|uv^T\|_F = \left( \sum_{i,j=1}^n u_i^2 v_j^2 \right)^{\frac{1}{2}} = \left( \sum_{i=1}^n u_i^2 \sum_{j=1}^n v_j^2 \right)^{\frac{1}{2}} = \|u\| \|v\|$$

**Proof.** (4/4).

Let  $B'$  and  $B''$  two different minimum. Then  $\frac{1}{2}(B' + B'') \in \mathcal{B}$  moreover

$$\left\| A - \frac{1}{2}(B' + B'') \right\|_F \leq \frac{1}{2} \|A - B'\|_F + \frac{1}{2} \|A - B''\|_F$$

If the inequality is strict we have a contradiction. From the Cauchy-Schwartz inequality we have an equality only when  $A - B' = \lambda(A - B'')$  so that

$$B' - \lambda B'' = (1 - \lambda)A$$

and

$$B's - \lambda B''s = (1 - \lambda)As \Rightarrow (1 - \lambda)y = (1 - \lambda)As$$

but this is true only when  $\lambda = 1$ , i.e.  $B' = B''$ .  $\square$

- 1 The update

$$A_{k+1} = A_k + \frac{(y_k - A_k s_k) s_k^T}{s_k^T s_k}$$

satisfy the secant condition:  $A_{k+1} s_k = y_k$  and  $A_{k+1}$  is the **nearest** matrix in the Frobenius norm that satisfy the secant condition.

- 2 Changing the norm we can have different results and in general you can loose uniqueness of the update.

## The Broyden method

(1/2)

### Algorithm (The Broyden method)

```

k ← 0; x0 and A0 assigned;
f0 ← F(x0);
while ‖fk‖ > ε do
  Solve for sk the linear system Aksk + fk = 0;
  xk+1 ← xk + sk;
  fk+1 ← F(xk+1);
  yk ← fk+1 - fk;
  Update: Ak+1 ← Ak +  $\frac{(y_k - A_k s_k) s_k^T}{s_k^T s_k}$ ;
  k ← k + 1;
end while

```

## The Brodyen method

(2/2)

Notice that  $\mathbf{y}_k - \mathbf{A}_k \mathbf{s}_k = \mathbf{f}_{k+1} - \mathbf{f}_k + \mathbf{f}_k$  so that the update can be written as  $\mathbf{A}_{k+1} \leftarrow \mathbf{A}_k + \mathbf{f}_{k+1} \mathbf{s}_k^T / \mathbf{s}_k^T \mathbf{s}_k$  and  $\mathbf{y}_k$  can be eliminated.

## Algorithm (The Brodyen method (alternative version))

```

k ← 0; x and A assigned;
f ← F(x);
while ||f|| > ε do
  Solve for s the linear system As + f = 0;
  x ← x + s;
  f ← F(x);
  Update: A ← A +  $\frac{\mathbf{f} \mathbf{s}^T}{\mathbf{s}^T \mathbf{s}}$ ;
  k ← k + 1;
end while
  
```



## Brodyen algorithm properties

(2/2)

## Theorem

Let  $\mathbf{F}(\mathbf{x}) = \mathbf{A}\mathbf{x} - \mathbf{b}$  where  $\mathbf{A} \in \mathbb{R}^{n \times n}$ . Then the Brodyen method converge in at most  $2n$  steps.

## Theorem

Let  $\mathbf{F} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  satisfy the standard regularity conditions with  $\nabla \mathbf{F}(\mathbf{x}_*)$  nonsingular. Then there exists positive constants  $\epsilon, \delta$  such that if  $\|\mathbf{x}_0 - \mathbf{x}_*\| \leq \epsilon$  and  $\|\mathbf{A}_0 - \nabla \mathbf{F}(\mathbf{x}_*)\| \leq \delta$ , then the sequence  $\{\mathbf{x}_k\}$  generated by the Brodyen method satisfy

$$\|\mathbf{x}_{k+2n} - \mathbf{x}_*\| \leq C \|\mathbf{x}_k - \mathbf{x}_*\|^2$$



D.M.Gay

Some convergence properties of Brodyen's method.  
SIAM J. Numer. Anal., 16 623-630, 1979.



## Brodyen algorithm properties

(1/2)

## Theorem

Let  $\mathbf{F}(\mathbf{x})$  satisfy the standard regularity conditions with  $\nabla \mathbf{F}(\mathbf{x}_*)$  nonsingular. Then there exists positive constants  $\epsilon, \delta$  such that if  $\|\mathbf{x}_0 - \mathbf{x}_*\| \leq \epsilon$  and  $\|\mathbf{A}_0 - \nabla \mathbf{F}(\mathbf{x}_*)\| \leq \delta$ , then the sequence  $\{\mathbf{x}_k\}$  generated by the Brodyen method is well defined and converge  $q$ -superlinearly to  $\mathbf{x}_*$ , i.e.

$$\lim_{k \rightarrow \infty} \frac{\|\mathbf{x}_{k+1} - \mathbf{x}_*\|}{\|\mathbf{x}_k - \mathbf{x}_*\|} = 0$$



C.G.Brodyen, J.E.Dennis, J.J.Moré

On the local and super-linear convergence of quasi-Newton methods.

J. Inst. Math. Appl, 6 222-236, 1973.



## Reorganizing Brodyen update

- Brodyen method needs to solve a linear system for  $\mathbf{A}_k$  at each step
- This can be onerous in terms of CPU cost
- it is possible to update directly the inverse of  $\mathbf{A}_k$  i.e. it is possible to update  $\mathbf{H}_k = \mathbf{A}_k^{-1}$ .
- The update of  $\mathbf{A}_k$  solve the problem of efficiency but do not alleviate the memory occupation
- The matrix  $\mathbf{A}_k$  can be written as a product of simple matrix, this can save memory if the update are lesser respect to the system dimension.



## Sherman-Morrison formula

Sherman-Morrison formula permit to explicit write the inverse of a matrix changed with a rank 1 perturbation

## Proposition (Sherman-Morrison formula)

$$(A + uv^T)^{-1} = A^{-1} - \frac{1}{\alpha} A^{-1} uv^T A^{-1}$$

where

$$\alpha = 1 + v^T A^{-1} u$$

The Sherman-Morrison formula can be checked by a direct calculation.



## Application of Sherman-Morrison formula

(2/2)

- The update formula for  $H_k$ :

$$H_{k+1} = H_k - \frac{1}{\beta_k} H_k f_{k+1} s_k^T H_k$$

$$\beta_k = s_k^T s_k + s_k^T H_k f_{k+1}$$

- Can be reorganized as follows

- Compute  $z_{k+1} = H_k f_{k+1}$ ;
- Compute  $\beta_k = s_k^T s_k + s_k^T z_{k+1}$ ;
- Compute  $H_{k+1} = (I - \beta_k^{-1} z_{k+1} s_k^T) H_k$ ;



## Application of Sherman-Morrison formula

(1/2)

- From the Broyden update formula

$$A_{k+1} = A_k + \frac{f_{k+1} s_k^T}{s_k^T s_k}$$

- By using Sherman-Morrison formula

$$A_{k+1}^{-1} = A_k^{-1} - \frac{1}{\beta_k} A_k^{-1} f_{k+1} s_k^T A_k^{-1}$$

$$\beta_k = s_k^T s_k + s_k^T A_k^{-1} f_{k+1}$$

- By setting  $H_k = A_k^{-1}$  we have the update formula for  $H_k$ :

$$H_{k+1} = H_k - \frac{1}{\beta_k} H_k f_{k+1} s_k^T H_k$$

$$\beta_k = s_k^T s_k + s_k^T H_k f_{k+1}$$



## The Broyden method with inverse updated

## Algorithm (The Broyden method (updating inverse))

```

k ← 0; x0 assigned;
f0 ← F(x0);
H0 ← I or better H0 ← ∇F(x0)-1;
while ||fk|| > ε do
  — perform step
  sk ← -Hk fk;
  xk+1 ← xk + sk;
  fk+1 ← F(xk+1);
  — update H
  zk+1 ← Hk fk+1;
  βk ← skT sk + skT zk+1;
  Hk+1 ← (I - βk-1 zk+1 skT) Hk;
  k ← k + 1;
end while

```



- If  $n$  is very large then the storing of  $H_k$  can be very expensive.
- Moreover when  $n$  is very large we hope to find a good solution with a number  $m$  of iteration with  $m \lll n$
- So that instead of storing  $H_k$  we can decide to store the vectors  $z_k$  and  $s_k$  plus the scalars  $\beta_k$ . With this vectors and scalars we can write

$$H_k = (I - \beta_{k-1} z_k s_{k-1}^T) \cdots (I - \beta_1 z_2 s_1^T) (I - \beta_0 z_1 s_0^T) H_0$$

- Assuming  $H_0 = I$  or can be computed on the fly we must store only  $2nm + m$  real number instead of  $n^2$  saving a lot of memory.
- However we can do better. It is possible to eliminate  $z_k$  and store only  $nm + m$  real numbers.

Elimination of  $z_k$ 

(2/3)

$$\begin{aligned} d_{k+1} &= H_{k+1} f_{k+1} = \left( I + \frac{z_{k+1} d_k^T}{d_k^T d_k - d_k^T z_{k+1}} \right) H_k f_{k+1} \\ &= \left( I + \frac{z_{k+1} d_k^T}{d_k^T d_k - d_k^T z_{k+1}} \right) z_{k+1} \\ &= z_{k+1} + \frac{z_{k+1} d_k^T z_{k+1}}{d_k^T d_k - d_k^T z_{k+1}} \\ &= \frac{d_k^T d_k}{d_k^T d_k - d_k^T z_{k+1}} z_{k+1} \end{aligned}$$

substituting in the update formula for  $H_{k+1}$  we obtain

$$H_{k+1} \leftarrow \left( I + \frac{d_{k+1} d_k^T}{d_k^T d_k} \right) H_k$$

Elimination of  $z_k$ 

(1/3)

- A step of the broyden iterative scheme can be rewritten as

$$d_k \leftarrow H_k f_k$$

$$x_{k+1} \leftarrow x_k - d_k$$

$$f_{k+1} \leftarrow F(x_{k+1})$$

$$z_{k+1} \leftarrow H_k f_{k+1}$$

$$H_{k+1} \leftarrow \left( I + \frac{z_{k+1} d_k^T}{d_k^T d_k - d_k^T z_{k+1}} \right) H_k$$

- you can notice that  $z_k$  and  $d_k$  are similar and contains a lot of common information.
- It is possible exploring the iteration to eliminate  $z_k$  from the update formula of  $H_k$  so that we can store the whole sequence without the vectors  $z_k$ .

Elimination of  $z_k$ 

(3/3)

Substituting into the step of the broyden iterative scheme and assuming  $d_k$  known

$$x_{k+1} \leftarrow x_k - d_k$$

$$f_{k+1} \leftarrow F(x_{k+1})$$

$$z_{k+1} \leftarrow H_k f_{k+1}$$

$$d_{k+1} \leftarrow \frac{d_k^T d_k}{d_k^T d_k - d_k^T z_{k+1}} z_{k+1}$$

$$H_{k+1} \leftarrow \left( I + \frac{d_{k+1} d_k^T}{d_k^T d_k} \right) H_k$$

notice that  $x_{k+1}$ ,  $f_{k+1}$  and  $z_{k+1}$  are not used in  $H_{k+1}$  so that only  $d_k$  and its length need to be stored.



## Algorithm (The Broyden method (low memory usage))

```

k ← 0; x assigned;
f ← F(x); H0 ← ∇F(x)-1; d0 ← H0f; ℓ0 ← d0Td0;
while ‖f‖ > ε do
  — perform step
  x ← x - dk;
  f ← F(x);
  — evaluate Hkf
  z ← H0f;
  for j = 0, 1, ..., k - 1 do
    z ← z + [(djTz)/ℓj]dj+1;
  end for
  — update Hk+1
  dk+1 ← [ℓk/(ℓk - dkTz)]z;
  ℓk+1 ← dk+1Tdk+1;
  k ← k + 1;
end while

```



## Outline

- 1 The Newton Raphson
- 2 The Broyden method
- 3 The dumped Broyden method



## Algorithm (The dumped Broyden method)

```

k ← 0; x0 assigned;
f0 ← F(x0); H0 ← ∇F(x0)-1;
while ‖fk‖ > ε do
  — compute search direction
  dk ← Hkfk;
  Approximate arg minλ>0 ‖F(xk - λdk)‖2 by line-search;
  — perform step
  sk ← -λkdk;
  xk+1 ← xk + sk;
  fk+1 ← F(xk+1);
  yk ← fk+1 - fk;
  — update Hk+1
  Hk+1 ← Hk + (sk - Hkyk)skT/skTHkyk;
  k ← k + 1;
end while

```

Elimination of z<sub>k</sub>

(1/5)

Notice that

$$H_k y_k = H_k f_{k+1} - H_k f_k = z_{k+1} - d_k, \quad \text{and} \quad s_k = -\lambda_k d_k$$

and

$$\begin{aligned}
 H_{k+1} &\leftarrow H_k + \frac{(s_k - H_k y_k) s_k^T}{s_k^T H_k y_k} H_k \\
 &\leftarrow H_k + \frac{(-\lambda_k d_k - z_{k+1} + d_k)(-\lambda_k d_k^T)}{-\lambda_k d_k^T (z_{k+1} - d_k)} H_k \\
 &\leftarrow \left( I + \frac{(-\lambda_k d_k - z_{k+1} + d_k) d_k^T}{d_k^T (z_{k+1} - d_k)} \right) H_k \\
 &\leftarrow \left( I + \frac{(z_{k+1} + (\lambda_k - 1) d_k) d_k^T}{d_k^T d_k - d_k^T z_{k+1}} \right) H_k
 \end{aligned}$$



Elimination of  $z_k$ 

(2/5)

A step of the brodyen iterative scheme can be rewritten as

$$\begin{aligned}d_k &\leftarrow H_k f_k \\x_{k+1} &\leftarrow x_k - \lambda_k d_k \\f_{k+1} &\leftarrow F(x_{k+1}) \\z_{k+1} &\leftarrow H_k f_{k+1} \\H_{k+1} &\leftarrow \left( I + \frac{(z_{k+1} + (\lambda_k - 1)d_k)d_k^T}{d_k^T d_k - d_k^T z_{k+1}} \right) H_k\end{aligned}$$

Elimination of  $z_k$ 

(3/5)

$$\begin{aligned}d_{k+1} &= H_{k+1} f_{k+1} \\&= \left( I + \frac{(z_{k+1} + (\lambda_k - 1)d_k)d_k^T}{d_k^T d_k - d_k^T z_{k+1}} \right) H_k f_{k+1} \\&= \left( I + \frac{(z_{k+1} + (\lambda_k - 1)d_k)d_k^T}{d_k^T d_k - d_k^T z_{k+1}} \right) z_{k+1} \\&= z_{k+1} + \frac{(z_{k+1} + (\lambda_k - 1)d_k)d_k^T z_{k+1}}{d_k^T d_k - d_k^T z_{k+1}} \\&= \frac{(d_k^T d_k)z_{k+1} + (\lambda_k - 1)(d_k^T z_{k+1})d_k}{d_k^T d_k - d_k^T z_{k+1}}\end{aligned}$$

Elimination of  $z_k$ 

(4/5)

Solving for  $z_{k+1}$

$$z_{k+1} = \frac{(d_k^T d_k - d_k^T z_{k+1})d_{k+1} - (\lambda_k - 1)(d_k^T z_{k+1})d_k}{d_k^T d_k}$$

and substituting in  $H_{k+1}$  we have

$$\begin{aligned}H_{k+1} &\leftarrow \left( I + \frac{(z_{k+1} + (\lambda_k - 1)d_k)d_k^T}{d_k^T d_k - d_k^T z_{k+1}} \right) H_k \\&\leftarrow \left( I + \frac{(d_{k+1} + (\lambda_k - 1)d_k)d_k^T}{d_k^T d_k} \right) H_k\end{aligned}$$

Elimination of  $z_k$ 

(5/5)

Substituting into the step of the brodyen iterative scheme and assuming  $d_k$  known

$$\begin{aligned}x_{k+1} &\leftarrow x_k - \lambda_k d_k \\f_{k+1} &\leftarrow F(x_{k+1}) \\z_{k+1} &\leftarrow H_k f_{k+1} \\d_{k+1} &\leftarrow \frac{(d_k^T d_k)z_{k+1} + (\lambda_k - 1)(d_k^T z_{k+1})d_k}{d_k^T d_k - d_k^T z_{k+1}} \\H_{k+1} &\leftarrow \left( I + \frac{(d_{k+1} + (\lambda_k - 1)d_k)d_k^T}{d_k^T d_k} \right) H_k\end{aligned}$$

notice that  $x_{k+1}$ ,  $f_{k+1}$  and  $z_{k+1}$  are not used in  $H_{k+1}$  so that only  $d_k$  and its length need to be stored.



## Algorithm (The dumped Broyden method)



```

 $k \leftarrow 0$ ;  $x$  assigned;
 $f \leftarrow F(x)$ ;  $H_0 \leftarrow \nabla F(x)^{-1}$ ;  $d_0 \leftarrow H_0 f$ ;  $\ell_0 \leftarrow d_0^T d_0$ ;
while  $\|f_k\| > \epsilon$  do
  Approximate  $\arg \min_{\lambda > 0} \|F(x - \lambda d_k)\|^2$  by line-search;
  — perform step
   $x \leftarrow x - \lambda_k d_k$ ;  $f \leftarrow F(x)$ ;
  — evaluate  $H_k f$ 
   $z \leftarrow H_0 f$ ;
  for  $j = 0, 1, \dots, k-1$  do
     $z \leftarrow z + [(d_j^T z)/\ell_j] (d_{j+1} + (\lambda_j - 1)d_j)$ ;
  end for
  — update  $H_{k+1}$ 
   $d_{k+1} \leftarrow [\ell_k z + (\lambda_k - 1)(d_k^T z)d_k] / (\ell_k - d_k^T z)$ ;
   $\ell_{k+1} \leftarrow d_{k+1}^T d_{k+1}$ ;
   $k \leftarrow k + 1$ ;
end while

```



## References

-  J. Stoer and R. Bulirsch  
 Introduction to numerical analysis  
 Springer-Verlag, Texts in Applied Mathematics, 12, 2002.
-  J. E. Dennis, Jr. and Robert B. Schnabel  
 Numerical Methods for Unconstrained Optimization and  
 Nonlinear Equations  
 SIAM, Classics in Applied Mathematics, 16, 1996.

