

Conjugate Direction minimization

Lectures for PHD course on
Unconstrained Numerical Optimization

Enrico Bertolazzi

DIMS – Università di Trento

May 2008



Outline

- 1 The Steepest Descent iterative scheme
- 2 Conjugate direction method
- 3 Conjugate Gradient method
- 4 Conjugate Gradient convergence rate
- 5 Preconditioning the Conjugate Gradient method
- 6 Nonlinear Conjugate Gradient extension



Generic minimization algorithm

In the following we study the convergence rate of the Generic minimization algorithm applied to a quadratic function $q(\mathbf{x})$ with **exact** line search. The function

$$q(\mathbf{x}) = \frac{1}{2} \mathbf{x}^T \mathbf{A} \mathbf{x} - \mathbf{b}^T \mathbf{x} + c$$

can be viewed as a n -dimensional generalization of the 1-dimensional parabolic model.

Generic minimization algorithm

```
Given an initial guess  $\mathbf{x}_0$ , let  $k = 0$ ;  
while not converged do  
  Find a descent direction  $\mathbf{p}_k$  at  $\mathbf{x}_k$ ;  
  Compute a step size  $\alpha_k$  using a line-search along  $\mathbf{p}_k$ .  
  Set  $\mathbf{x}_{k+1} = \mathbf{x}_k + \alpha_k \mathbf{p}_k$  and increase  $k$  by 1.  
end while
```



Assumption (Symmetry)

The matrix \mathbf{A} is assumed to be symmetric, in fact,

$$\mathbf{A} = \mathbf{A}^{Symm} + \mathbf{A}^{Skew}$$

where

$$\mathbf{A}^{Symm} = \frac{1}{2} [\mathbf{A} + \mathbf{A}^T], \quad \mathbf{A}^{Symm} = (\mathbf{A}^{Symm})^T$$

$$\mathbf{A}^{Skew} = \frac{1}{2} [\mathbf{A} - \mathbf{A}^T], \quad \mathbf{A}^{Skew} = -(\mathbf{A}^{Skew})^T$$

moreover

$$\mathbf{x}^T \mathbf{A} \mathbf{x} = \mathbf{x}^T \mathbf{A}^{Symm} \mathbf{x} + \mathbf{x}^T \mathbf{A}^{Skew} \mathbf{x} = \mathbf{x}^T \mathbf{A}^{Symm} \mathbf{x}$$

so that only the symmetric part of \mathbf{A} contribute to $q(\mathbf{x})$.



Assumption (SPD)

The matrix \mathbf{A} is assumed to be symmetric and positive definite, in fact,

$$\nabla q(\mathbf{x})^T = \frac{1}{2}(\mathbf{A} + \mathbf{A}^T)\mathbf{x} - \mathbf{b} = \mathbf{A}\mathbf{x} - \mathbf{b}$$

and

$$\nabla^2 q(\mathbf{x}) = \frac{1}{2}(\mathbf{A} + \mathbf{A}^T) = \mathbf{A}$$

From the *sufficient* condition for a minimum we have that $\nabla q(\mathbf{x}_\star)^T = \mathbf{0}$, i.e.

$$\mathbf{A}\mathbf{x}_\star = \mathbf{b}$$

and $\nabla^2 q(\mathbf{x}_\star) = \mathbf{A}$ is SPD.



The toy problem

(1/3)

- In the following we study the convergence rate of the Steepest Descent and Conjugate Gradient methods applied to

$$q(\mathbf{x}) = \frac{1}{2}\mathbf{x}^T \mathbf{A}\mathbf{x} - \mathbf{b}^T \mathbf{x} + c$$

where \mathbf{A} is an SPD matrix.

- This assumption simplify the analysis but it is also useful in the non linear case. In fact, by expanding a generic function $f(\mathbf{x})$ near its minimum \mathbf{x}_\star we have

$$\begin{aligned} f(\mathbf{x}) &= f(\mathbf{x}_\star) + \nabla f(\mathbf{x}_\star)(\mathbf{x} - \mathbf{x}_\star) \\ &\quad + \frac{1}{2}(\mathbf{x} - \mathbf{x}_\star)^T \nabla^2 f(\mathbf{x}_\star)(\mathbf{x} - \mathbf{x}_\star) + \mathcal{O}(\|\mathbf{x} - \mathbf{x}_\star\|^3) \end{aligned}$$



- By setting

$$\mathbf{A} = \nabla^2 f(\mathbf{x}_*),$$

$$\mathbf{b} = \nabla^2 f(\mathbf{x}_*)\mathbf{x}_* - \nabla f(\mathbf{x}_*)$$

$$c = f(\mathbf{x}_*) - \nabla f(\mathbf{x}_*)\mathbf{x}_* + \frac{1}{2}\mathbf{x}_*^T \nabla^2 f(\mathbf{x}_*)\mathbf{x}_*$$

we have

$$f(\mathbf{x}) = \frac{1}{2}\mathbf{x}^T \mathbf{A}\mathbf{x} - \mathbf{b}^T \mathbf{x} + c + \mathcal{O}(\|\mathbf{x} - \mathbf{x}_*\|^3)$$

- So that we expect that when an iterate \mathbf{x}_k is near \mathbf{x}_* then we can neglect $\mathcal{O}(\|\mathbf{x} - \mathbf{x}_*\|^3)$ and the asymptotic behavior is the same of the quadratic problem.



- we can rewrite the quadratic problem in many different way as follows

$$\begin{aligned} q(\mathbf{x}) &= \frac{1}{2}(\mathbf{x} - \mathbf{x}_*)^T \mathbf{A}(\mathbf{x} - \mathbf{x}_*) + c' \\ &= \frac{1}{2}(\mathbf{A}\mathbf{x} - \mathbf{b})^T \mathbf{A}^{-1}(\mathbf{A}\mathbf{x} - \mathbf{b}) + c' \end{aligned}$$

where

$$c' = c + \frac{1}{2}\mathbf{x}_*^T \mathbf{A}\mathbf{x}_*$$

- This last forms are useful in the study of the steepest descent method.



Outline

- 1 The Steepest Descent iterative scheme
- 2 Conjugate direction method
- 3 Conjugate Gradient method
- 4 Conjugate Gradient convergence rate
- 5 Preconditioning the Conjugate Gradient method
- 6 Nonlinear Conjugate Gradient extension

The steepest descent for quadratic functions

(1/3)

The steepest descent minimization algorithm

Given an initial guess \mathbf{x}_0 , let $k = 0$;

while not converged do

 Choose as descent direction $\mathbf{p}_k = -\nabla q(\mathbf{x}_k)^T = \mathbf{b} - \mathbf{A}\mathbf{x}_k$;

 Compute a step size α_k using a line-search along \mathbf{p}_k .

 Set $\mathbf{x}_{k+1} = \mathbf{x}_k + \alpha_k \mathbf{p}_k$ and increase k by 1.

end while

Definition (Residual)

The expressions

$$\mathbf{r}(\mathbf{x}) = \mathbf{b} - \mathbf{A}\mathbf{x}, \quad \mathbf{r}_k = \mathbf{b} - \mathbf{A}\mathbf{x}_k$$

are called the residual. We obviously have $\mathbf{r}(\mathbf{x}) = -\nabla q(\mathbf{x})^T$ and $\mathbf{r}(\mathbf{x}_) = \mathbf{0}$.*

The steepest descent for quadratic functions

(2/3)

Lemma

The solution of the minimization problem:

$$\alpha_k = \arg \min_{\alpha \geq 0} q(\mathbf{x}_k - \alpha \mathbf{r}_k) \quad \text{is} \quad \alpha_k = -\frac{\mathbf{r}_k^T \mathbf{r}_k}{\mathbf{r}_k^T \mathbf{A} \mathbf{r}_k}.$$

Proof.

Because $p(\alpha) = q(\mathbf{x}_k - \alpha \mathbf{r}_k)$ the minimum is a stationary point:

$$\begin{aligned} \frac{dp(\alpha)}{d\alpha} &= \frac{dq(\mathbf{x}_k - \alpha \mathbf{r}_k)}{d\alpha} = -\nabla q(\mathbf{x}_k - \alpha \mathbf{r}_k) \mathbf{r}_k \\ &= \mathbf{r}(\mathbf{x}_k - \alpha \mathbf{r}_k)^T \mathbf{r}_k = (\mathbf{b} - \mathbf{A}(\mathbf{x}_k - \alpha \mathbf{r}_k))^T \mathbf{r}_k \\ &= (\mathbf{r}_k + \alpha \mathbf{A} \mathbf{r}_k)^T \mathbf{r}_k = 0 \end{aligned}$$

and solving for α the result follows. □



The steepest descent for quadratic functions

(3/3)

The steepest descent minimization algorithm

Given an initial guess \mathbf{x}_0 , let $k = 0$;

while not converged do

 Compute $\mathbf{r}_k = \mathbf{b} - \mathbf{A} \mathbf{x}_k$;

 Compute the step size $\alpha_k = \frac{\mathbf{r}_k^T \mathbf{r}_k}{\mathbf{r}_k^T \mathbf{A} \mathbf{r}_k}$;

 Set $\mathbf{x}_{k+1} = \mathbf{x}_k + \alpha_k \mathbf{r}_k$ and increase k by 1.

end while

Or more compactly

$$\mathbf{x}_{k+1} = \mathbf{x}_k + \frac{\mathbf{r}_k^T \mathbf{r}_k}{\mathbf{r}_k^T \mathbf{A} \mathbf{r}_k} \mathbf{r}_k$$



The steepest descent reduction step

(1/4)

The next lemma bound the reduction of $q(\mathbf{x}_{k+1})$ by the value of $q(\mathbf{x}_k)$:

Lemma

Consider the steepest descent for quadratic function, than we have the following estimate

$$\|\mathbf{x}_* - \mathbf{x}_{k+1}\|_A^2 = \|\mathbf{x}_* - \mathbf{x}_k\|_A^2 \left(1 - \frac{(\mathbf{r}_k^T \mathbf{r}_k)^2}{(\mathbf{r}_k^T \mathbf{A}^{-1} \mathbf{r}_k)(\mathbf{r}_k^T \mathbf{A} \mathbf{r}_k)} \right)$$

where

$$\|\mathbf{x}\|_A = \sqrt{\mathbf{x}^T \mathbf{A} \mathbf{x}}$$

is the **energy norm** induced by the SPD matrix \mathbf{A} .



The steepest descent reduction step

(2/4)

Proof.

(1/3).

We want bound $q(\mathbf{x}_{k+1})$ by $q(\mathbf{x}_k)$:

$$\begin{aligned} q(\mathbf{x}_{k+1}) &= q(\mathbf{x}_k + \alpha_k \mathbf{r}_k) \\ &= \frac{1}{2} (\mathbf{A} \mathbf{x}_k + \alpha_k \mathbf{A} \mathbf{r}_k - \mathbf{b})^T \mathbf{A}^{-1} (\mathbf{A} \mathbf{x}_k + \alpha_k \mathbf{A} \mathbf{r}_k - \mathbf{b}) + c' \\ &= \frac{1}{2} (\alpha_k \mathbf{A} \mathbf{r}_k - \mathbf{r}_k)^T \mathbf{A}^{-1} (\alpha_k \mathbf{A} \mathbf{r}_k - \mathbf{r}_k) + c' \\ &= \frac{1}{2} \mathbf{r}_k^T \mathbf{A}^{-1} \mathbf{r}_k + \frac{1}{2} \alpha_k^2 \mathbf{r}_k^T \mathbf{A} \mathbf{r}_k - \alpha_k \mathbf{r}_k^T \mathbf{r}_k + c' \\ &= q(\mathbf{x}_k) + \frac{1}{2} \alpha_k (\alpha_k \mathbf{r}_k^T \mathbf{A} \mathbf{r}_k - 2 \mathbf{r}_k^T \mathbf{r}_k) \end{aligned}$$



The steepest descent reduction step

(3/4)

Proof.

(2/3).

Substituting $\alpha_k = \frac{\mathbf{r}_k^T \mathbf{r}_k}{\mathbf{r}_k^T \mathbf{A} \mathbf{r}_k}$ we obtain

$$q(\mathbf{x}_{k+1}) = q(\mathbf{x}_k) - \frac{1}{2} \frac{(\mathbf{r}_k^T \mathbf{r}_k)^2}{\mathbf{r}_k^T \mathbf{A} \mathbf{r}_k}$$

this shows that the steepest descent method reduce at each step the objective function $q(\mathbf{x})$.

Using the expression $q(\mathbf{x}) = \frac{1}{2} \mathbf{r}(\mathbf{x})^T \mathbf{A}^{-1} \mathbf{r}(\mathbf{x}) + c'$ we can write:

$$\frac{1}{2} \mathbf{r}_{k+1}^T \mathbf{A}^{-1} \mathbf{r}_{k+1} = \frac{1}{2} \mathbf{r}_k^T \mathbf{A}^{-1} \mathbf{r}_k - \frac{1}{2} \frac{(\mathbf{r}_k^T \mathbf{r}_k)^2}{\mathbf{r}_k^T \mathbf{A} \mathbf{r}_k}$$



The steepest descent reduction step

(4/4)

Proof.

(3/3).

or better

$$\mathbf{r}_{k+1}^T \mathbf{A}^{-1} \mathbf{r}_{k+1} = \mathbf{r}_k^T \mathbf{A}^{-1} \mathbf{r}_k \left(1 - \frac{(\mathbf{r}_k^T \mathbf{r}_k)^2}{(\mathbf{r}_k^T \mathbf{A}^{-1} \mathbf{r}_k)(\mathbf{r}_k^T \mathbf{A} \mathbf{r}_k)} \right)$$

noticing that $\mathbf{r}_k = \mathbf{b} - \mathbf{A} \mathbf{x}_k = \mathbf{A} \mathbf{x}_* - \mathbf{A} \mathbf{x}_k = \mathbf{A}(\mathbf{x}_* - \mathbf{x}_k)$ we have

$$\|\mathbf{x}_* - \mathbf{x}_{k+1}\|_{\mathbf{A}}^2 = \|\mathbf{x}_* - \mathbf{x}_k\|_{\mathbf{A}}^2 \left(1 - \frac{(\mathbf{r}_k^T \mathbf{r}_k)^2}{(\mathbf{r}_k^T \mathbf{A}^{-1} \mathbf{r}_k)(\mathbf{r}_k^T \mathbf{A} \mathbf{r}_k)} \right)$$

where

$$\|\mathbf{x}\|_{\mathbf{A}} = \sqrt{\mathbf{x}^T \mathbf{A} \mathbf{x}}$$

is the **energy norm** induced by the SPD matrix \mathbf{A} .



The estimate of the convergence rate for the **steepest descent** method is linked to the estimate of the term

$$\frac{(\mathbf{r}_k^T \mathbf{r}_k)^2}{(\mathbf{r}_k^T \mathbf{A}^{-1} \mathbf{r}_k)(\mathbf{r}_k^T \mathbf{A} \mathbf{r}_k)}$$

in particular we can prove

Lemma (Kantorovic)

Let $\mathbf{A} \in \mathbb{R}^{n \times n}$ an SPD matrix then the following inequality is valid

$$1 \leq \frac{(\mathbf{x}^T \mathbf{A} \mathbf{x})(\mathbf{x}^T \mathbf{A}^{-1} \mathbf{x})}{(\mathbf{x}^T \mathbf{x})^2} \leq \frac{(M + m)^2}{4 M m}$$

for all $\mathbf{x} \neq \mathbf{0}$. Where $m = \lambda_1$ is the smallest eigenvalue of \mathbf{A} and $M = \lambda_n$ is the biggest eigenvalue of \mathbf{A} .



Proof.

(1/5).

STEP 1: problem reformulation. First of all notice that

$$\frac{(\mathbf{x}^T \mathbf{A} \mathbf{x})(\mathbf{x}^T \mathbf{A}^{-1} \mathbf{x})}{(\mathbf{x}^T \mathbf{x})^2} = \frac{(\mathbf{y}^T \mathbf{A} \mathbf{y})(\mathbf{y}^T \mathbf{A}^{-1} \mathbf{y})}{(\mathbf{y}^T \mathbf{y})^2}$$

for all $\mathbf{y} = \alpha \mathbf{x}$ with $\alpha \neq 0$. Choosing $\alpha = \|\mathbf{x}\|^{-1}$ have:

$$\min_{\|\mathbf{z}\|=1} (\mathbf{z}^T \mathbf{A} \mathbf{z})(\mathbf{z}^T \mathbf{A}^{-1} \mathbf{z}) \leq$$

$$\frac{(\mathbf{x}^T \mathbf{A} \mathbf{x})(\mathbf{x}^T \mathbf{A}^{-1} \mathbf{x})}{(\mathbf{x}^T \mathbf{x})^2}$$

$$\leq \max_{\|\mathbf{z}\|=1} (\mathbf{z}^T \mathbf{A} \mathbf{z})(\mathbf{z}^T \mathbf{A}^{-1} \mathbf{z})$$



Proof.

(2/5).

STEP 2: eigenvector expansions. Matrix $\mathbf{A} \in \mathbb{R}^{n \times n}$ is an SPD matrix so that there exists $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n$ a complete orthonormal eigenvectors set with $0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$ corresponding eigenvalues. Let be $\mathbf{x} \in \mathbb{R}^n$ then

$$\mathbf{x} = \sum_{k=1}^n \alpha_k \mathbf{u}_k, \quad \mathbf{x}^T \mathbf{x} = \sum_{k=1}^n \alpha_k^2$$

so that $(\mathbf{x}^T \mathbf{A} \mathbf{x})(\mathbf{x}^T \mathbf{A}^{-1} \mathbf{x}) = h(\alpha_1, \dots, \alpha_n)$ where

$$h(\alpha_1, \dots, \alpha_n) = \left(\sum_{k=1}^n \alpha_k^2 \lambda_k \right) \left(\sum_{k=1}^n \alpha_k^2 \lambda_k^{-1} \right)$$

then the lemma can be reformulated:

- Find maxima and minima of $h(\alpha_1, \dots, \alpha_n)$
- subject to $\sum_{k=1}^n \alpha_k^2 = 1$.



Proof.

(3/5).

STEP 3: problem reduction. By using Lagrange multiplier maxima and minima are the stationary points of:

$$g(\alpha_1, \dots, \alpha_n, \mu) = h(\alpha_1, \dots, \alpha_n) + \mu \left(\sum_{k=1}^n \alpha_k^2 - 1 \right)$$

setting $A = \sum_{k=1}^n \alpha_k^2 \lambda_k$ and $B = \sum_{k=1}^n \alpha_k^2 \lambda_k^{-1}$ we have

$$\frac{\partial g(\alpha_1, \dots, \alpha_n, \mu)}{\partial \alpha_k} = 2\alpha_k (\lambda_k B + \lambda_k^{-1} A + \mu) = 0$$

so that

- 1 Or $\alpha_k = 0$;
- 2 Or λ_k is a root of the quadratic polynomial $\lambda^2 B + \lambda \mu + A$.

in any case there are at most 2 coefficients α 's not zero. ^a

^athe argument should be improved in the case of multiple eigenvalues



Proof.

(4/5).

STEP 4: problem reformulation. say α_i and α_j are the only non zero coefficients, then $\alpha_i^2 + \alpha_j^2 = 1$ and we can write

$$\begin{aligned}
 h(\alpha_1, \dots, \alpha_n) &= (\alpha_i^2 \lambda_i + \alpha_j^2 \lambda_j) (\alpha_i^2 \lambda_i^{-1} + \alpha_j^2 \lambda_j^{-1}) \\
 &= \alpha_i^4 + \alpha_j^4 + \alpha_i^2 \alpha_j^2 \left(\frac{\lambda_i}{\lambda_j} + \frac{\lambda_j}{\lambda_i} \right) \\
 &= \alpha_i^2 (1 - \alpha_j^2) + \alpha_j^2 (1 - \alpha_i^2) + \alpha_i^2 \alpha_j^2 \left(\frac{\lambda_i}{\lambda_j} + \frac{\lambda_j}{\lambda_i} \right) \\
 &= 1 + \alpha_i^2 \alpha_j^2 \left(\frac{\lambda_i}{\lambda_j} + \frac{\lambda_j}{\lambda_i} - 2 \right) \\
 &= 1 + \alpha_i^2 (1 - \alpha_i^2) \frac{(\lambda_i - \lambda_j)^2}{\lambda_i \lambda_j}
 \end{aligned}$$



Proof.

(5/5).

STEP 5: bounding maxima and minima. notice that

$$0 \leq \beta(1 - \beta) \leq \frac{1}{4}, \quad \forall \beta \in [0, 1]$$

$$1 \leq 1 + \alpha_i^2 (1 - \alpha_i^2) \frac{(\lambda_i - \lambda_j)^2}{\lambda_i \lambda_j} \leq 1 + \frac{(\lambda_i - \lambda_j)^2}{4 \lambda_i \lambda_j} = \frac{(\lambda_i + \lambda_j)^2}{4 \lambda_i \lambda_j}$$

to bound $(\lambda_i + \lambda_j)^2 / (4 \lambda_i \lambda_j)$ consider the function $f(x) = (1 + x)^2 / x$ which is increasing for $x \geq 1$ so that we have

$$\frac{(\lambda_i + \lambda_j)^2}{4 \lambda_i \lambda_j} \leq \frac{(M + m)^2}{4 M m}$$

and finally

$$1 \leq h(\alpha_1, \dots, \alpha_n) \leq \frac{(M + m)^2}{4 M m}$$



Convergence rate of Steepest Descent

The Kantorovich inequality permits to prove:

Theorem (Convergence rate of Steepest Descent)

Let $\mathbf{A} \in \mathbb{R}^{n \times n}$ an SPD matrix then the **steepest descent** method:

$$\mathbf{x}_{k+1} = \mathbf{x}_k + \frac{\mathbf{r}_k^T \mathbf{r}_k}{\mathbf{r}_k^T \mathbf{A} \mathbf{r}_k} \mathbf{r}_k$$

converge to the solution $\mathbf{x}_* = \mathbf{A}^{-1} \mathbf{b}$ with at least linear q -rate in the norm $\|\cdot\|_{\mathbf{A}}$. Moreover we have the error estimate

$$\|\mathbf{x}_{k+1} - \mathbf{x}_*\|_{\mathbf{A}} \leq \frac{\kappa - 1}{\kappa + 1} \|\mathbf{x}_k - \mathbf{x}_*\|_{\mathbf{A}}$$

$\kappa = M/m$ is the **condition number** where $m = \lambda_1$ is the smallest eigenvalue of \mathbf{A} and $M = \lambda_n$ is the biggest eigenvalue of \mathbf{A} .



Proof.

Remember from slide $N^{\circ}16$

$$\|\mathbf{x}_* - \mathbf{x}_{k+1}\|_{\mathbf{A}}^2 = \|\mathbf{x}_* - \mathbf{x}_k\|_{\mathbf{A}}^2 \left(1 - \frac{(\mathbf{r}_k^T \mathbf{r}_k)^2}{(\mathbf{r}_k^T \mathbf{A}^{-1} \mathbf{r}_k)(\mathbf{r}_k^T \mathbf{A} \mathbf{r}_k)} \right)$$

from Kantorovich inequality

$$1 - \frac{(\mathbf{r}_k^T \mathbf{r}_k)^2}{(\mathbf{r}_k^T \mathbf{A}^{-1} \mathbf{r}_k)(\mathbf{r}_k^T \mathbf{A} \mathbf{r}_k)} \leq 1 - \frac{4 M m}{(M + m)^2} = \frac{(M - m)^2}{(M + m)^2}$$

so that

$$\|\mathbf{x}_* - \mathbf{x}_{k+1}\|_{\mathbf{A}} \leq \frac{M - m}{M + m} \|\mathbf{x}_* - \mathbf{x}_k\|_{\mathbf{A}}$$

□



Remark (One step convergence)

The steepest descent method can converge in one iteration if $\kappa = 1$ or when $\mathbf{r}_0 = \mathbf{u}_k$ where \mathbf{u}_k is an eigenvector of \mathbf{A} .

- 1 In the first case ($\kappa = 1$) we have $\mathbf{A} = \beta \mathbf{I}$ for some $\beta > 0$ so it is not interesting.
- 2 In the second case we have

$$\frac{(\mathbf{u}_k^T \mathbf{u}_k)^2}{(\mathbf{u}_k^T \mathbf{A}^{-1} \mathbf{u}_k)(\mathbf{u}_k^T \mathbf{A} \mathbf{u}_k)} = \frac{(\mathbf{u}_k^T \mathbf{u}_k)^2}{\lambda_k^{-1}(\mathbf{u}_k^T \mathbf{u}_k) \lambda_k(\mathbf{u}_k^T \mathbf{u}_k)} = 1$$

in both cases we have $\mathbf{r}_1 = \mathbf{0}$ i.e. we have found the solution.



Outline

- 1 The Steepest Descent iterative scheme
- 2 Conjugate direction method
- 3 Conjugate Gradient method
- 4 Conjugate Gradient convergence rate
- 5 Preconditioning the Conjugate Gradient method
- 6 Nonlinear Conjugate Gradient extension



Conjugate direction method

Definition (Conjugate vector)

Given two vectors \mathbf{p} and \mathbf{q} in \mathbb{R}^n are **conjugate respect to \mathbf{A}** if they are orthogonal respect the scalar product induced by \mathbf{A} ; i.e.,

$$\mathbf{p}^T \mathbf{A} \mathbf{q} = \sum_{i,j=1}^n A_{ij} p_i q_j = 0.$$

Clearly, n vectors $\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_n \in \mathbb{R}^n$ that are pair wise conjugated respect to \mathbf{A} form a base of \mathbb{R}^n .



Problem (Linear system)

Find the minimum of $q(\mathbf{x}) = \frac{1}{2} \mathbf{x}^T \mathbf{A} \mathbf{x} - \mathbf{b}^T \mathbf{x} + c$ is equivalent to solve the first order necessary condition, i.e.

$$\text{Find } \mathbf{x}_* \in \mathbb{R}^n \text{ such that: } \mathbf{A} \mathbf{x}_* = \mathbf{b}.$$

Observation

Consider $\mathbf{x}_0 \in \mathbb{R}^n$ and decompose the error $\mathbf{e}_0 = \mathbf{x}_* - \mathbf{x}_0$ by the conjugate vectors $\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_n \in \mathbb{R}^n$:

$$\mathbf{e}_0 = \mathbf{x}_* - \mathbf{x}_0 = \sigma_1 \mathbf{p}_1 + \sigma_2 \mathbf{p}_2 + \dots + \sigma_n \mathbf{p}_n.$$

Evaluating the coefficients $\sigma_1, \sigma_2, \dots, \sigma_n \in \mathbb{R}$ is equivalent to solve the problem $\mathbf{A} \mathbf{x}_* = \mathbf{b}$, because knowing \mathbf{e}_0 we have

$$\mathbf{x}_* = \mathbf{x}_0 + \mathbf{e}_0.$$



Observation

Using conjugacy the coefficients $\sigma_1, \sigma_2, \dots, \sigma_n \in \mathbb{R}$ can be computed as

$$\sigma_i = \frac{\mathbf{p}_i^T \mathbf{A} \mathbf{e}_0}{\mathbf{p}_i^T \mathbf{A} \mathbf{p}_i}, \quad \text{for } i = 1, 2, \dots, n.$$

In fact, for all $1 \leq i \leq n$, we have

$$\begin{aligned} \mathbf{p}_i^T \mathbf{A} \mathbf{e}_0 &= \mathbf{p}_i^T \mathbf{A} (\sigma_1 \mathbf{p}_1 + \sigma_2 \mathbf{p}_2 + \dots + \sigma_n \mathbf{p}_n), \\ &= \sigma_1 \mathbf{p}_i^T \mathbf{A} \mathbf{p}_1 + \sigma_2 \mathbf{p}_i^T \mathbf{A} \mathbf{p}_2 + \dots + \sigma_n \mathbf{p}_i^T \mathbf{A} \mathbf{p}_n, \\ &= \sigma_i \mathbf{p}_i^T \mathbf{A} \mathbf{p}_i, \end{aligned}$$

because $\mathbf{p}_i^T \mathbf{A} \mathbf{p}_j = 0$ for $i \neq j$.



The conjugate direction method evaluate the coefficients $\sigma_1, \sigma_2, \dots, \sigma_n \in \mathbb{R}$ recursively in n steps, solving for $k \geq 0$ the minimization problem:

Conjugate direction method

Given \mathbf{x}_0 ; $k \leftarrow 0$;

repeat

$k \leftarrow k + 1$;

Find $\mathbf{x}_k \in \mathbf{x}_0 + \mathcal{V}_k$ such that:

$$\mathbf{x}_k = \arg \min_{\mathbf{x} \in \mathbf{x}_0 + \mathcal{V}_k} \|\mathbf{x}_* - \mathbf{x}\|_{\mathbf{A}}$$

until $k = n$

where \mathcal{V}_k is the subspace of \mathbb{R}^n generated by the first k conjugate direction; i.e.,

$$\mathcal{V}_k = \text{SPAN}\{\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_k\}.$$



Step: $\mathbf{x}_0 \rightarrow \mathbf{x}_1$

At the first step we consider the subspace $\mathbf{x}_0 + \text{SPAN}\{\mathbf{p}_1\}$ which consists in vectors of the form

$$\mathbf{x}(\alpha) = \mathbf{x}_0 + \alpha\mathbf{p}_1 \quad \alpha \in \mathbb{R}$$

The minimization problem becomes:

Minimization step $\mathbf{x}_0 \rightarrow \mathbf{x}_1$

Find $\mathbf{x}_1 = \mathbf{x}_0 + \alpha_1\mathbf{p}_1$ (i.e., find $\alpha_1!$) such that:

$$\|\mathbf{x}_* - \mathbf{x}_1\|_{\mathbf{A}} = \min_{\alpha \in \mathbb{R}} \|\mathbf{x}_* - (\mathbf{x}_0 + \alpha\mathbf{p}_1)\|_{\mathbf{A}},$$



Solving first step method 1

The minimization problem is the minimum respect to α of the quadratic:

$$\begin{aligned} \Phi(\alpha) &= \|\mathbf{x}_* - (\mathbf{x}_0 + \alpha\mathbf{p}_1)\|_{\mathbf{A}}^2, \\ &= (\mathbf{x}_* - (\mathbf{x}_0 + \alpha\mathbf{p}_1))^T \mathbf{A} (\mathbf{x}_* - (\mathbf{x}_0 + \alpha\mathbf{p}_1)), \\ &= (\mathbf{e}_0 - \alpha\mathbf{p}_1)^T \mathbf{A} (\mathbf{e}_0 - \alpha\mathbf{p}_1), \\ &= \mathbf{e}_0^T \mathbf{A} \mathbf{e}_0 - 2\alpha \mathbf{p}_1^T \mathbf{A} \mathbf{e}_0 + \alpha^2 \mathbf{p}_1^T \mathbf{A} \mathbf{p}_1. \end{aligned}$$

minimum is found by imposing:

$$\frac{d\Phi(\alpha)}{d\alpha} = -2\mathbf{p}_1^T \mathbf{A} \mathbf{e}_0 + 2\alpha \mathbf{p}_1^T \mathbf{A} \mathbf{p}_1 = 0 \quad \Rightarrow$$

$$\alpha_1 = \frac{\mathbf{p}_1^T \mathbf{A} \mathbf{e}_0}{\mathbf{p}_1^T \mathbf{A} \mathbf{p}_1}$$



Solving first step method 2

(1/2)

Remember the error expansion:

$$\mathbf{x}_* - \mathbf{x}_0 = \sigma_1 \mathbf{p}_1 + \sigma_2 \mathbf{p}_2 + \cdots + \sigma_n \mathbf{p}_n.$$

Let $\mathbf{x}(\alpha) = \mathbf{x}_0 + \alpha \mathbf{p}_1$, the difference $\mathbf{x}_* - \mathbf{x}(\alpha)$ becomes:

$$\mathbf{x}_* - \mathbf{x}(\alpha) = (\sigma_1 - \alpha) \mathbf{p}_1 + \sigma_2 \mathbf{p}_2 + \cdots + \sigma_n \mathbf{p}_n$$

due to conjugacy the error $\|\mathbf{x}_* - \mathbf{x}(\alpha)\|_{\mathbf{A}}$ becomes

$$\begin{aligned} \|\mathbf{x}_* - \mathbf{x}(\alpha)\|_{\mathbf{A}}^2 &= \left((\sigma_1 - \alpha) \mathbf{p}_1 + \sum_{i=2}^n \sigma_i \mathbf{p}_i \right)^T \mathbf{A} \left((\sigma_1 - \alpha) \mathbf{p}_1 + \sum_{j=2}^n \sigma_j \mathbf{p}_j \right) \\ &= (\sigma_1 - \alpha)^2 \mathbf{p}_1^T \mathbf{A} \mathbf{p}_1 + \sum_{j=2}^n \sigma_j^2 \mathbf{p}_j^T \mathbf{A} \mathbf{p}_j \end{aligned}$$



Solving first step method 2

(2/2)

Because

$$\|\mathbf{x}_* - \mathbf{x}(\alpha)\|_{\mathbf{A}}^2 = (\sigma_1 - \alpha)^2 \|\mathbf{p}_1\|_{\mathbf{A}}^2 + \sum_{i=2}^n \sigma_i^2 \|\mathbf{p}_i\|_{\mathbf{A}}^2,$$

we have that

$$\|\mathbf{x}_* - \mathbf{x}(\alpha_1)\|_{\mathbf{A}}^2 = \sum_{i=2}^n \sigma_i^2 \|\mathbf{p}_i\|_{\mathbf{A}}^2 \leq \|\mathbf{x}_* - \mathbf{x}(\alpha)\|_{\mathbf{A}}^2 \quad \text{for all } \alpha \neq \sigma_1$$

so that minimum is found by imposing $\alpha_1 = \sigma_1$:

$$\alpha_1 = \frac{\mathbf{p}_1^T \mathbf{A} \mathbf{e}_0}{\mathbf{p}_1^T \mathbf{A} \mathbf{p}_1}$$

This argument can be generalized for all $k > 1$ (see next slides).



Step, $\mathbf{x}_{k-1} \rightarrow \mathbf{x}_k$

For the step from $k - 1$ to k we consider the subspace of \mathbb{R}^n

$$\mathcal{V}_k = \text{SPAN}\{\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_k\}$$

which contains vectors of the form:

$$\mathbf{x}(\alpha^{(1)}, \alpha^{(2)}, \dots, \alpha^{(k)}) = \mathbf{x}_0 + \alpha^{(1)}\mathbf{p}_1 + \alpha^{(2)}\mathbf{p}_2 + \dots + \alpha^{(k)}\mathbf{p}_k$$

The minimization problem becomes:

Minimization step $\mathbf{x}_{k-1} \rightarrow \mathbf{x}_k$

Find $\mathbf{x}_k = \mathbf{x}_0 + \alpha_1\mathbf{p}_1 + \alpha_2\mathbf{p}_2 + \dots + \alpha_k\mathbf{p}_k$ (i.e. $\alpha_1, \alpha_2, \dots, \alpha_k$) such that:

$$\|\mathbf{x}_\star - \mathbf{x}_k\|_{\mathbf{A}} = \min_{\alpha^{(1)}, \alpha^{(2)}, \dots, \alpha^{(k)} \in \mathbb{R}} \left\| \mathbf{x}_\star - \mathbf{x}(\alpha^{(1)}, \alpha^{(2)}, \dots, \alpha^{(k)}) \right\|_{\mathbf{A}}$$

Solving kth Step: $\mathbf{x}_{k-1} \rightarrow \mathbf{x}_k$

(1/2)

Remember the error expansion:

$$\mathbf{x}_\star - \mathbf{x}_0 = \sigma_1\mathbf{p}_1 + \sigma_2\mathbf{p}_2 + \dots + \sigma_n\mathbf{p}_n.$$

Consider a vector of the form

$$\mathbf{x}(\alpha^{(1)}, \alpha^{(2)}, \dots, \alpha^{(k)}) = \mathbf{x}_0 + \alpha^{(1)}\mathbf{p}_1 + \alpha^{(2)}\mathbf{p}_2 + \dots + \alpha^{(k)}\mathbf{p}_k$$

the error $\mathbf{x}_\star - \mathbf{x}(\alpha^{(1)}, \alpha^{(2)}, \dots, \alpha^{(k)})$ can be written as

$$\begin{aligned} \mathbf{x}_\star - \mathbf{x}(\alpha^{(1)}, \alpha^{(2)}, \dots, \alpha^{(k)}) &= \mathbf{x}_\star - \mathbf{x}_0 - \sum_{i=1}^k \alpha^{(i)}\mathbf{p}_i, \\ &= \sum_{i=1}^k (\sigma_i - \alpha^{(i)})\mathbf{p}_i + \sum_{i=k+1}^n \sigma_i\mathbf{p}_i. \end{aligned}$$



Solving k th Step: $\mathbf{x}_{k-1} \rightarrow \mathbf{x}_k$

(2/2)

using conjugacy of \mathbf{p}_i we obtain the norm of the error:

$$\begin{aligned} & \left\| \mathbf{x}_* - \mathbf{x}(\alpha^{(1)}, \alpha^{(2)}, \dots, \alpha^{(k)}) \right\|_{\mathbf{A}}^2 \\ &= \sum_{i=1}^k (\sigma_i - \alpha^{(i)})^2 \|\mathbf{p}_i\|_{\mathbf{A}}^2 + \sum_{i=k+1}^n \sigma_i^2 \|\mathbf{p}_i\|_{\mathbf{A}}^2. \end{aligned}$$

So that minimum is found by imposing $\alpha_i = \sigma_i$: for $i = 1, 2, \dots, k$.

$$\alpha_i = \frac{\mathbf{p}_i^T \mathbf{A} \mathbf{e}_0}{\mathbf{p}_i^T \mathbf{A} \mathbf{p}_i} \quad i = 1, 2, \dots, k$$



Successive one dimensional minimization

(1/3)

- notice that $\alpha_i = \sigma_i$ and that

$$\begin{aligned} \mathbf{x}_k &= \mathbf{x}_0 + \alpha_1 \mathbf{p}_1 + \dots + \alpha_k \mathbf{p}_k \\ &= \mathbf{x}_{k-1} + \alpha_k \mathbf{p}_k \end{aligned}$$

- so that \mathbf{x}_{k-1} contains $k - 1$ coefficients α_i for the minimization.
- if we consider the one dimensional minimization on the subspace $\mathbf{x}_{k-1} + \text{SPAN}\{\mathbf{p}_k\}$ we find again \mathbf{x}_k !



Successive one dimensional minimization

(2/3)

Consider a vector of the form

$$\mathbf{x}(\alpha) = \mathbf{x}_{k-1} + \alpha \mathbf{p}_k$$

remember that $\mathbf{x}_{k-1} = \mathbf{x}_0 + \alpha_1 \mathbf{p}_1 + \dots + \alpha_{k-1} \mathbf{p}_{k-1}$ so that the error $\mathbf{x}_* - \mathbf{x}(\alpha)$ can be written as

$$\begin{aligned} \mathbf{x}_* - \mathbf{x}(\alpha) &= \mathbf{x}_* - \mathbf{x}_0 - \sum_{i=1}^{k-1} \alpha_i \mathbf{p}_i + \alpha \mathbf{p}_k \\ &= \sum_{i=1}^{k-1} (\sigma_i - \alpha_i) \mathbf{p}_i + (\sigma_k - \alpha) \mathbf{p}_k + \sum_{i=k+1}^n \sigma_i \mathbf{p}_i. \end{aligned}$$

due to the equality $\sigma_i = \alpha_i$ the blue part of the expression is 0.



Successive one dimensional minimization

(3/3)

Using conjugacy of \mathbf{p}_i we obtain the norm of the error:

$$\|\mathbf{x}_* - \mathbf{x}(\alpha)\|_{\mathbf{A}}^2 = (\sigma_k - \alpha)^2 \|\mathbf{p}_k\|_{\mathbf{A}}^2 + \sum_{i=k+1}^n \sigma_i^2 \|\mathbf{p}_i\|_{\mathbf{A}}^2.$$

So that minimum is found by imposing $\alpha = \sigma_k$:

$$\alpha_k = \frac{\mathbf{p}_k^T \mathbf{A} \mathbf{e}_0}{\mathbf{p}_k^T \mathbf{A} \mathbf{p}_k}$$

Remark

This observation permit to perform the minimization on the k -dimensional space $\mathbf{x}_0 + \mathcal{V}_k$ as successive one dimensional minimizations along the conjugate directions \mathbf{p}_k !



Problem (one dimensional successive minimization)

Find $\mathbf{x}_k = \mathbf{x}_{k-1} + \alpha_k \mathbf{p}_k$ such that:

$$\|\mathbf{x}_* - \mathbf{x}_k\|_{\mathbf{A}} = \min_{\alpha \in \mathbb{R}} \|\mathbf{x}_* - (\mathbf{x}_{k-1} + \alpha \mathbf{p}_k)\|_{\mathbf{A}},$$

The solution is the minimum respect to α of the quadratic:

$$\begin{aligned} \Phi(\alpha) &= (\mathbf{x}_* - (\mathbf{x}_{k-1} + \alpha \mathbf{p}_k))^T \mathbf{A} (\mathbf{x}_* - (\mathbf{x}_{k-1} + \alpha \mathbf{p}_k)), \\ &= (\mathbf{e}_{k-1} - \alpha \mathbf{p}_k)^T \mathbf{A} (\mathbf{e}_{k-1} - \alpha \mathbf{p}_k), \\ &= \mathbf{e}_{k-1}^T \mathbf{A} \mathbf{e}_{k-1} - 2\alpha \mathbf{p}_k^T \mathbf{A} \mathbf{e}_{k-1} + \alpha^2 \mathbf{p}_k^T \mathbf{A} \mathbf{p}_k. \end{aligned}$$

minimum is found by imposing:

$$\frac{d\Phi(\alpha)}{d\alpha} = -2\mathbf{p}_k^T \mathbf{A} \mathbf{e}_{k-1} + 2\alpha \mathbf{p}_k^T \mathbf{A} \mathbf{p}_k = 0 \quad \Rightarrow$$

$$\alpha_k = \frac{\mathbf{p}_k^T \mathbf{A} \mathbf{e}_{k-1}}{\mathbf{p}_k^T \mathbf{A} \mathbf{p}_k}$$



- In the case of minimization on the subspace $\mathbf{x}_0 + \mathcal{V}_k$ we have:

$$\alpha_k = \mathbf{p}_k^T \mathbf{A} \mathbf{e}_0 / \mathbf{p}_k^T \mathbf{A} \mathbf{p}_k$$

- In the case of one dimensional minimization on the subspace $\mathbf{x}_{k-1} + \text{SPAN}\{\mathbf{p}_k\}$ we have:

$$\alpha_k = \mathbf{p}_k^T \mathbf{A} \mathbf{e}_{k-1} / \mathbf{p}_k^T \mathbf{A} \mathbf{p}_k$$

- Apparently they are different results, however by using the conjugacy of the vectors \mathbf{p}_i we have

$$\begin{aligned} \mathbf{p}_k^T \mathbf{A} \mathbf{e}_{k-1} &= \mathbf{p}_k^T \mathbf{A} (\mathbf{x}_* - \mathbf{x}_{k-1}) \\ &= \mathbf{p}_k^T \mathbf{A} (\mathbf{x}_* - (\mathbf{x}_0 + \alpha_1 \mathbf{p}_1 + \dots + \alpha_{k-1} \mathbf{p}_{k-1})) \\ &= \mathbf{p}_k^T \mathbf{A} \mathbf{e}_0 - \alpha_1 \mathbf{p}_k^T \mathbf{A} \mathbf{p}_1 - \dots - \alpha_{k-1} \mathbf{p}_k^T \mathbf{A} \mathbf{p}_{k-1} \\ &= \mathbf{p}_k^T \mathbf{A} \mathbf{e}_0 \end{aligned}$$



- The **one step minimization** in the space $\mathbf{x}_0 + \mathcal{V}_n$ and the **successive minimization** in the space $\mathbf{x}_{k-1} + \text{SPAN}\{\mathbf{p}_k\}$, $k = 1, 2, \dots, n$ are equivalent if \mathbf{p}_i s are conjugate.
- The successive minimization is useful when \mathbf{p}_i s are not known in advance but must be computed as the minimization process proceeds.
- The evaluation of α_k is apparently not computable because \mathbf{e}_i is not known. However noticing

$$\mathbf{A}\mathbf{e}_k = \mathbf{A}(\mathbf{x}_* - \mathbf{x}_k) = \mathbf{b} - \mathbf{A}\mathbf{x}_k = \mathbf{r}_k$$

we can write

$$\alpha_k = \mathbf{p}_k^T \mathbf{A}\mathbf{e}_{k-1} / \mathbf{p}_k^T \mathbf{A}\mathbf{p}_k = \mathbf{p}_k^T \mathbf{r}_{k-1} / \mathbf{p}_k^T \mathbf{A}\mathbf{p}_k =$$

- Finally for the residual is valid the recurrence

$$\mathbf{r}_k = \mathbf{b} - \mathbf{A}\mathbf{x}_k = \mathbf{b} - \mathbf{A}(\mathbf{x}_{k-1} + \alpha_k \mathbf{p}_k) = \mathbf{r}_{k-1} - \alpha_k \mathbf{A}\mathbf{p}_k.$$



Conjugate direction minimization

Algorithm (Conjugate direction minimization)

```

k ← 0; x0 assigned;
r0 ← b - Ax0;
while not converged do
  k ← k + 1;
  αk ← (rk-1T pkT) / (pkT Apk);
  xk ← xk-1 + αk pk;
  rk ← rk-1 - αk Apk;
end while

```

Observation (Computational cost)

The conjugate direction minimization requires at each step one matrix–vector product for the evaluation of α_k and two update **AXPY** for \mathbf{x}_k and \mathbf{r}_k .



Monotonic behavior of the error

Remark (Monotonic behavior of the error)

The **energy norm** of the error $\|e_k\|_A$ is monotonically decreasing in k . In fact:

$$e_k = \mathbf{x}_* - \mathbf{x}_k = \alpha_{k+1}\mathbf{p}_{k+1} + \dots + \alpha_n\mathbf{p}_n,$$

and by conjugacy

$$\|e_k\|_A^2 = \|\mathbf{x}_* - \mathbf{x}_k\|_A^2 = \sigma_{k+1}^2 \|\mathbf{p}_{k+1}\|_A^2 + \dots + \sigma_n^2 \|\mathbf{p}_n\|_A^2.$$

Finally from this relation we have $e_n = \mathbf{0}$.



Outline

- 1 The Steepest Descent iterative scheme
- 2 Conjugate direction method
- 3 Conjugate Gradient method**
- 4 Conjugate Gradient convergence rate
- 5 Preconditioning the Conjugate Gradient method
- 6 Nonlinear Conjugate Gradient extension



Conjugate Gradient method

The Conjugate Gradient method combine the **Conjugate Direction** method with an **orthogonalization process** (like Gram-Schmidt) applied to the residual to construct the conjugate directions. In fact, because \mathbf{A} define a scalar product in the next slide we prove:

- each residue is orthogonal to the previous conjugate directions, and consequently linearly independent from the previous conjugate directions.
- if the residual is not null is can be used to construct a new conjugate direction.



Orthogonality of the residue \mathbf{r}_k respect \mathcal{V}_k

- The residue \mathbf{r}_k is orthogonal to $\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_k$. In fact, from the error expansion

$$\mathbf{e}_k = \alpha_{k+1}\mathbf{p}_{k+1} + \alpha_{k+2}\mathbf{p}_{k+2} + \dots + \alpha_n\mathbf{p}_n$$

because $\mathbf{r}_k = \mathbf{A}\mathbf{e}_k$, for $i = 1, 2, \dots, k$ we have

$$\begin{aligned} \mathbf{p}_i^T \mathbf{r}_k &= \mathbf{p}_i^T \mathbf{A}\mathbf{e}_k \\ &= \mathbf{p}_i^T \mathbf{A} \sum_{j=k+1}^n \alpha_j \mathbf{p}_j = \sum_{j=k+1}^n \alpha_j \mathbf{p}_i^T \mathbf{A}\mathbf{p}_j \\ &= 0 \end{aligned}$$



Building new conjugate direction

(1/2)

- The conjugate direction method build **one new** direction at each step.
- If $\mathbf{r}_k \neq \mathbf{0}$ it can be used to build the new direction \mathbf{p}_{k+1} by a Gram-Schmidt orthogonalization process

$$\mathbf{p}_{k+1} = \mathbf{r}_k + \beta_1^{(k+1)} \mathbf{p}_1 + \beta_2^{(k+1)} \mathbf{p}_2 + \dots + \beta_k^{(k+1)} \mathbf{p}_k,$$

where the k coefficients $\beta_1^{(k+1)}, \beta_2^{(k+1)}, \dots, \beta_k^{(k+1)}$ must satisfy:

$$\mathbf{p}_i^T \mathbf{A} \mathbf{p}_{k+1} = 0, \quad \text{for } i = 1, 2, \dots, k.$$



Building new conjugate direction

(2/2)

(repeating from previous slide)

$$\mathbf{p}_{k+1} = \mathbf{r}_k + \beta_1^{(k+1)} \mathbf{p}_1 + \beta_2^{(k+1)} \mathbf{p}_2 + \dots + \beta_k^{(k+1)} \mathbf{p}_k,$$

expanding the expression:

$$\begin{aligned} 0 &= \mathbf{p}_i^T \mathbf{A} \mathbf{p}_{k+1}, \\ &= \mathbf{p}_i^T \mathbf{A} (\mathbf{r}_k + \beta_1^{(k+1)} \mathbf{p}_1 + \beta_2^{(k+1)} \mathbf{p}_2 + \dots + \beta_k^{(k+1)} \mathbf{p}_k), \\ &= \mathbf{p}_i^T \mathbf{A} \mathbf{r}_k + \beta_i^{(k+1)} \mathbf{p}_i^T \mathbf{A} \mathbf{p}_i, \end{aligned}$$

$$\Rightarrow \boxed{\beta_i^{(k+1)} = -\frac{\mathbf{p}_i^T \mathbf{A} \mathbf{r}_k}{\mathbf{p}_i^T \mathbf{A} \mathbf{p}_i}} \quad i = 1, 2, \dots, k$$



The choice of the residual $\mathbf{r}_k \neq \mathbf{0}$ for the construction of the new conjugate direction \mathbf{p}_{k+1} has **three** important consequences:

- ① simplification of the expression for α_k ;
- ② Orthogonality of the residual \mathbf{r}_k from the previous residue $\mathbf{r}_0, \mathbf{r}_1, \dots, \mathbf{r}_{k-1}$;
- ③ **three point formula** and simplification of the coefficients $\beta_i^{(k+1)}$.

this facts will be examined in the next slides.

Simplification of the expression for α_k

Writing the expression for \mathbf{p}_k from the orthogonalization process

$$\mathbf{p}_k = \mathbf{r}_{k-1} + \beta_1^{(k+1)} \mathbf{p}_1 + \beta_2^{(k+1)} \mathbf{p}_2 + \dots + \beta_{k-1}^{(k+1)} \mathbf{p}_{k-1},$$

using orthogonality of \mathbf{r}_{k-1} and the vectors $\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_{k-1}$, (see slide N.48) we have

$$\begin{aligned} \mathbf{r}_{k-1}^T \mathbf{p}_k &= \mathbf{r}_{k-1}^T (\mathbf{r}_{k-1} + \beta_1^{(k+1)} \mathbf{p}_1 + \beta_2^{(k+1)} \mathbf{p}_2 + \dots + \beta_{k-1}^{(k+1)} \mathbf{p}_{k-1}), \\ &= \mathbf{r}_{k-1}^T \mathbf{r}_{k-1}. \end{aligned}$$

recalling the definition of α_k it follows:

$$\alpha_k = \frac{\mathbf{e}_{k-1}^T \mathbf{A} \mathbf{p}_k}{\mathbf{p}_k^T \mathbf{A} \mathbf{p}_k} = \frac{\mathbf{r}_{k-1}^T \mathbf{p}_k}{\mathbf{p}_k^T \mathbf{A} \mathbf{p}_k} = \boxed{\frac{\mathbf{r}_{k-1}^T \mathbf{r}_{k-1}}{\mathbf{p}_k^T \mathbf{A} \mathbf{p}_k}}$$

Orthogonality of the residue \mathbf{r}_k from $\mathbf{r}_0, \mathbf{r}_1, \dots, \mathbf{r}_{k-1}$

From the definition of \mathbf{p}_{i+1} it follows:

$$\mathbf{p}_{i+1} = \mathbf{r}_i + \beta_1^{(i+1)} \mathbf{p}_1 + \beta_2^{(i+1)} \mathbf{p}_2 + \dots + \beta_i^{(i+1)} \mathbf{p}_i,$$

$$\Rightarrow \mathbf{r}_i \in \text{SPAN}\{\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_i, \mathbf{p}_{i+1}\} = \mathcal{V}_{i+1} \quad (\text{obvious})$$

using orthogonality of \mathbf{r}_k and the vectors $\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_k$, (see slide N.48) for $i < k$ we have

$$\begin{aligned} \mathbf{r}_k^T \mathbf{r}_i &= \mathbf{r}_k^T \left(\mathbf{p}_{i+1} - \sum_{j=1}^i \beta_j^{(i+1)} \mathbf{p}_j \right), \\ &= \mathbf{r}_k^T \mathbf{p}_{i+1} - \sum_{j=1}^i \beta_j^{(i+1)} \mathbf{r}_k^T \mathbf{p}_j = 0. \end{aligned}$$



Three point formula and simplification of $\beta_i^{(k+1)}$

From the relation $\mathbf{r}_k^T \mathbf{r}_i = \mathbf{r}_k^T (\mathbf{r}_{i-1} - \alpha_i \mathbf{A} \mathbf{p}_i)$ we deduce

$$\mathbf{r}_k^T \mathbf{A} \mathbf{p}_i = \frac{\mathbf{r}_k^T \mathbf{r}_{i-1} - \mathbf{r}_k^T \mathbf{r}_i}{\alpha_i} = \begin{cases} -\mathbf{r}_k^T \mathbf{r}_k / \alpha_k & \text{if } i = k; \\ 0 & \text{if } i < k; \end{cases}$$

remembering that $\alpha_k = \mathbf{r}_{k-1}^T \mathbf{r}_{k-1} / \mathbf{p}_k^T \mathbf{A} \mathbf{p}_k$ we obtain

$$\beta_i^{(k+1)} = -\frac{\mathbf{r}_k^T \mathbf{A} \mathbf{p}_i}{\mathbf{p}_i^T \mathbf{A} \mathbf{p}_i} = \begin{cases} \frac{\mathbf{r}_k^T \mathbf{r}_k}{\mathbf{r}_{k-1}^T \mathbf{r}_{k-1}} & i = k; \\ 0 & i < k; \end{cases}$$

i.e. there is only one non zero coefficient $\beta_k^{(k+1)}$, so we write $\beta_k = \beta_k^{(k+1)}$ and obtain the **three point formula**:

$$\mathbf{p}_{k+1} = \mathbf{r}_k + \beta_k \mathbf{p}_k$$



Conjugate gradient algorithm

initial step:

$k \leftarrow 0$; \mathbf{x}_0 assigned;

$\mathbf{r}_0 \leftarrow \mathbf{b} - \mathbf{A}\mathbf{x}_0$;

$\mathbf{p}_1 \leftarrow \mathbf{r}_0$;

while $\|\mathbf{r}_k\| > \epsilon$ **do**

$k \leftarrow k + 1$;

Conjugate direction method

$$\alpha_k \leftarrow \frac{\mathbf{r}_{k-1}^T \mathbf{r}_{k-1}}{\mathbf{p}_k^T \mathbf{A} \mathbf{p}_k};$$

$$\mathbf{x}_k \leftarrow \mathbf{x}_{k-1} + \alpha_k \mathbf{p}_k;$$

$$\mathbf{r}_k \leftarrow \mathbf{r}_{k-1} - \alpha_k \mathbf{A} \mathbf{p}_k;$$

Residual orthogonalization

$$\beta_k \leftarrow \frac{\mathbf{r}_k^T \mathbf{r}_k}{\mathbf{r}_{k-1}^T \mathbf{r}_{k-1}};$$

$$\mathbf{p}_{k+1} \leftarrow \mathbf{r}_k + \beta_k \mathbf{p}_k;$$

end while

Outline

- 1 The Steepest Descent iterative scheme
- 2 Conjugate direction method
- 3 Conjugate Gradient method
- 4 Conjugate Gradient convergence rate
- 5 Preconditioning the Conjugate Gradient method
- 6 Nonlinear Conjugate Gradient extension

Polynomial residual expansions

(1/6)

Lemma

The residuals and conjugate directions for the Conjugate Gradient iterative scheme of slide 55 can be written as

$$\mathbf{r}_k = P_k(\mathbf{A})\mathbf{r}_0 \quad k = 0, 1, \dots, n$$

$$\mathbf{p}_k = Q_{k-1}(\mathbf{A})\mathbf{r}_0 \quad k = 1, 2, \dots, n$$

where $P_k(x)$ and $Q_k(x)$ are k -degree polynomial such that $P_k(0) = 1$ for all k .

Proof.

(1/2).

The proof is by induction.

Base $k = 0$: $\mathbf{p}_1 = \mathbf{r}_0$
so that $P_0(x) = 1$ and $Q_0(x) = 1$.



Polynomial residual expansions

(2/6)

Proof.

(2/2).

Let the expansion valid for $k - 1$. Consider the recursion for the residual:

$$\begin{aligned} \mathbf{r}_k &= \mathbf{r}_{k-1} - \alpha_k \mathbf{A} \mathbf{p}_k \\ &= P_{k-1}(\mathbf{A})\mathbf{r}_0 + \alpha_k \mathbf{A} Q_{k-1}(\mathbf{A})\mathbf{r}_0 \\ &= (P_{k-1}(\mathbf{A}) + \alpha_k \mathbf{A} Q_{k-1}(\mathbf{A}))\mathbf{r}_0 \end{aligned}$$

then $P_k(x) = P_{k-1}(x) + \alpha_k x Q_{k-1}(x)$ and $P_k(0) = P_{k-1}(0) = 1$. Consider the recursion for the conjugate direction

$$\begin{aligned} \mathbf{p}_{k+1} &= P_k(\mathbf{A})\mathbf{r}_0 + \beta_k Q_{k-1}(\mathbf{A})\mathbf{r}_0 \\ &= (P_k(\mathbf{A}) + \beta_k Q_{k-1}(\mathbf{A}))\mathbf{r}_0 \end{aligned}$$

then $Q_k(x) = P_k(x) + \beta_k Q_{k-1}(x)$. □



Polynomial residual expansions

(3/6)

Corollary

$$\mathbf{e}_k = P_k(\mathbf{A})\mathbf{e}_0.$$

Proof.

$$\begin{aligned} \mathbf{e}_k &= \mathbf{x}_\star - \mathbf{x}_k = \mathbf{A}^{-1}\mathbf{r}_k \\ &= \mathbf{A}^{-1}P_k(\mathbf{A})\mathbf{r}_0 \\ &= P_k(\mathbf{A})\mathbf{A}^{-1}\mathbf{r}_0 \\ &= P_k(\mathbf{A})(\mathbf{x}_\star - \mathbf{x}_0) \\ &= P_k(\mathbf{A})\mathbf{e}_0. \end{aligned}$$



Polynomial residual expansions

(4/6)

Lemma

For the Conjugate Gradient iterative scheme of slide n.55 we have:

$$\mathcal{V}_k = \{p(\mathbf{A})\mathbf{e}_0 \mid p \in \mathbb{P}^k, p(0) = 0\}$$

Proof.

Using expansion of slide n.57 and $\mathbf{r}_0 = \mathbf{A}\mathbf{e}_0$ we have:

$$\begin{aligned} \mathcal{V}_k &= \text{SPAN}\{\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_k\} \\ &= \left\{ \sum_{i=0}^{k-1} \beta_i Q_i(\mathbf{A})\mathbf{r}_0 \mid (\beta_0, \dots, \beta_{k-1}) \in \mathbb{R}^{k-1} \right\} \\ &= \{q(\mathbf{A})\mathbf{A}\mathbf{e}_0 \mid q \in \mathbb{P}^{k-1}\} = \{p(\mathbf{A})\mathbf{e}_0 \mid p \in \mathbb{P}^k, p(0) = 0\} \end{aligned}$$



Polynomial residual expansions

(5/6)

By using the equality

$$\mathcal{V}_k = \{p(\mathbf{A})\mathbf{e}_0 \mid p \in \mathbb{P}^k, p(0) = 0\}$$

The optimality of CG step can be written as

$$\|\mathbf{x}_* - \mathbf{x}_k\|_{\mathbf{A}} \leq \|\mathbf{x}_* - \mathbf{x}\|_{\mathbf{A}}, \quad \forall \mathbf{x} \in \mathbf{x}_0 + \mathcal{V}_k$$

$$\|\mathbf{x}_* - \mathbf{x}_k\|_{\mathbf{A}} \leq \|\mathbf{x}_* - (\mathbf{x}_0 + p(\mathbf{A})\mathbf{e}_0)\|_{\mathbf{A}}, \quad \forall p \in \mathbb{P}^k, p(0) = 0$$

$$\|\mathbf{x}_* - \mathbf{x}_k\|_{\mathbf{A}} \leq \|P(\mathbf{A})\mathbf{e}_0\|_{\mathbf{A}}, \quad \forall P \in \mathbb{P}^k, P(0) = 1$$

And using the results of slide 60 and 59 we can write

$$\mathbf{e}_k = P_k(\mathbf{A})\mathbf{e}_0,$$

$$\|\mathbf{e}_k\|_{\mathbf{A}} = \|P_k(\mathbf{A})\mathbf{e}_0\|_{\mathbf{A}} \leq \|P(\mathbf{A})\mathbf{e}_0\|_{\mathbf{A}} \quad \forall P \in \mathbb{P}^k, P(0) = 1$$



Polynomial residual expansions

(6/6)

From previous equations we have the characterization of CG error

$$\|\mathbf{e}_k\|_{\mathbf{A}} = \inf_{P \in \mathbb{P}^k, P(0)=1} \|P(\mathbf{A})\mathbf{e}_0\|_{\mathbf{A}}$$

Thus, an estimate of the form

$$\|\mathbf{e}_k\|_{\mathbf{A}} \leq C_k \|\mathbf{e}_0\|_{\mathbf{A}}$$

can be obtained by using estimate on the polynomial of the form

$$\{P \in \mathbb{P}^k, P(0) = 1\}$$



Convergence rate calculation

Lemma

Let $\mathbf{A} \in \mathbb{R}^{n \times n}$ an SPD matrix, and $p \in \mathbb{P}^k$ a polynomial, then

$$\|p(\mathbf{A})\mathbf{x}\|_{\mathbf{A}} \leq \|p(\mathbf{A})\|_2 \|\mathbf{x}\|_{\mathbf{A}}$$

Proof.

(1/2).

The matrix \mathbf{A} is SPD so that we can write

$$\mathbf{A} = \mathbf{U}^T \mathbf{\Lambda} \mathbf{U}, \quad \mathbf{\Lambda} = \text{DIAG}\{\lambda_1, \lambda_2, \dots, \lambda_n\}$$

where \mathbf{U} is an orthogonal matrix (i.e. $\mathbf{U}^T \mathbf{U} = \mathbf{I}$) and $\mathbf{\Lambda} \geq \mathbf{0}$ is diagonal. We can define the SPD matrix $\mathbf{A}^{1/2}$ as follows

$$\mathbf{A}^{1/2} = \mathbf{U}^T \mathbf{\Lambda}^{1/2} \mathbf{U}, \quad \mathbf{\Lambda}^{1/2} = \text{DIAG}\{\lambda_1^{1/2}, \lambda_2^{1/2}, \dots, \lambda_n^{1/2}\}$$

and obviously $\mathbf{A}^{1/2} \mathbf{A}^{1/2} = \mathbf{A}$.



Proof.

(2/2).

Notice that

$$\|\mathbf{x}\|_{\mathbf{A}}^2 = \mathbf{x}^T \mathbf{A} \mathbf{x} = \mathbf{x}^T \mathbf{A}^{1/2} \mathbf{A}^{1/2} \mathbf{x} = \left\| \mathbf{A}^{1/2} \mathbf{x} \right\|_2^2$$

so that

$$\begin{aligned} \|p(\mathbf{A})\mathbf{x}\|_{\mathbf{A}} &= \left\| \mathbf{A}^{1/2} p(\mathbf{A}) \mathbf{x} \right\|_2 \\ &= \left\| p(\mathbf{A}) \mathbf{A}^{1/2} \mathbf{x} \right\|_2 \\ &\leq \|p(\mathbf{A})\|_2 \left\| \mathbf{A}^{1/2} \mathbf{x} \right\|_2 \\ &= \|p(\mathbf{A})\|_2 \|\mathbf{x}\|_{\mathbf{A}} \end{aligned}$$



Lemma

Let $\mathbf{A} \in \mathbb{R}^{n \times n}$ an SPD matrix, and $p \in \mathbb{P}^k$ a polynomial, then

$$\|p(\mathbf{A})\|_2 = \max_{\lambda \in \sigma(\mathbf{A})} |p(\lambda)|$$

Proof.

The matrix $p(\mathbf{A})$ is symmetric, and for a generic symmetric matrix \mathbf{B} we have

$$\|\mathbf{B}\|_2 = \max_{\lambda \in \sigma(\mathbf{B})} |\lambda|$$

observing that if λ is an eigenvalue of \mathbf{A} then $p(\lambda)$ is an eigenvalue of $p(\mathbf{A})$ the thesis easily follows. \square

- Starting the error estimate

$$\|e_k\|_{\mathbf{A}} \leq \inf_{P \in \mathbb{P}^k, P(0)=1} \|P(\mathbf{A})e_0\|_{\mathbf{A}}$$

- Combining the last two lemma we easily obtain the estimate

$$\|e_k\|_{\mathbf{A}} \leq \inf_{P \in \mathbb{P}^k, P(0)=1} \left[\max_{\lambda \in \sigma(\mathbf{A})} |P(\lambda)| \right] \|e_0\|_{\mathbf{A}}$$

- The convergence rate is estimated by bounding the constant

$$\inf_{P \in \mathbb{P}^k, P(0)=1} \left[\max_{\lambda \in \sigma(\mathbf{A})} |P(\lambda)| \right]$$

Finite termination of Conjugate Gradient

Theorem (Finite termination of Conjugate Gradient)

Let $\mathbf{A} \in \mathbb{R}^{n \times n}$ an SPD matrix, the the **Conjugate Gradient** applied to the linear system $\mathbf{A}\mathbf{x} = \mathbf{b}$ terminate finding the exact solution in at most n -step.

Proof.

From the estimate

$$\|\mathbf{e}_k\|_{\mathbf{A}} \leq \inf_{P \in \mathbb{P}^k, P(0)=1} \left[\max_{\lambda \in \sigma(\mathbf{A})} |P(\lambda)| \right] \|\mathbf{e}_0\|_{\mathbf{A}}$$

choosing
$$P(x) = \prod_{\lambda \in \sigma(\mathbf{A})} (x - \lambda) / \prod_{\lambda \in \sigma(\mathbf{A})} (0 - \lambda)$$

we have $\max_{\lambda \in \sigma(\mathbf{A})} |P(\lambda)| = 0$ and $\|\mathbf{e}_n\|_{\mathbf{A}} = 0$. □

Convergence rate of Conjugate Gradient

- 1 The constant

$$\inf_{P \in \mathbb{P}^k, P(0)=1} \left[\max_{\lambda \in \sigma(\mathbf{A})} |P(\lambda)| \right]$$

is not easy to evaluate,

- 2 The following bound, is useful

$$\max_{\lambda \in \sigma(\mathbf{A})} |P(\lambda)| \leq \max_{\lambda \in [\lambda_1, \lambda_n]} |P(\lambda)|$$

- 3 in particular the final estimate will be obtained by

$$\inf_{P \in \mathbb{P}^k, P(0)=1} \left[\max_{\lambda \in \sigma(\mathbf{A})} |P(\lambda)| \right] \leq \max_{\lambda \in [\lambda_1, \lambda_n]} |\bar{P}_k(\lambda)|$$

where $\bar{P}_k(x)$ is an opportune k -degree polynomial for which $\bar{P}_k(0) = 1$ and it is easy to evaluate $\max_{\lambda \in [\lambda_1, \lambda_n]} |\bar{P}_k(\lambda)|$.

Chebyshev Polynomials

(1/4)

- 1 The **Chebyshev Polynomials of the First Kind** are the right polynomial for this estimate. This polynomial have the following definition in the interval $[-1, 1]$:

$$T_k(x) = \cos(k \arccos(x))$$

- 2 Another equivalent definition valid in the interval $(-\infty, \infty)$ is the following

$$T_k(x) = \frac{1}{2} \left[\left(x + \sqrt{x^2 - 1} \right)^k + \left(x - \sqrt{x^2 - 1} \right)^k \right]$$





- 3 In spite of these definition, $T_k(x)$ is effectively a polynomial.



Chebyshev Polynomials

(2/4)

Some example of Chebyshev Polynomials.



Chebyshev Polynomials

(3/4)

- ① It is easy to show that $T_k(x)$ is a polynomial by the use of

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

$$\cos(\alpha + \beta) + \cos(\alpha - \beta) = 2 \cos \alpha \cos \beta$$

let $\theta = \arccos(x)$:

- ① $T_0(x) = \cos(0\theta) = 1$;
- ② $T_1(x) = \cos(1\theta) = x$;
- ③ $T_2(x) = \cos(2\theta) = \cos^2(\theta) - \sin^2(\theta) = 2\cos^2(\theta) - 1 = 2x^2 - 1$;
- ④ $T_{k+1}(x) + T_{k-1}(x) = \cos((k+1)\theta) + \cos((k-1)\theta)$
 $= 2\cos(k\theta)\cos(\theta) = 2xT_k(x)$

- ② In general we have the following recurrence:

- ① $T_0(x) = 1$;
- ② $T_1(x) = x$;
- ③ $T_{k+1}(x) = 2xT_k(x) - T_{k-1}(x)$.



Chebyshev Polynomials

(4/4)

- Solving the recurrence:
 - ① $T_0(x) = 1$;
 - ② $T_1(x) = x$;
 - ③ $T_{k+1}(x) = 2xT_k(x) - T_{k-1}(x)$.
- We obtain the explicit form of the Chebyshev Polynomials

$$T_k(x) = \frac{1}{2} \left[\left(x + \sqrt{x^2 - 1} \right)^k + \left(x - \sqrt{x^2 - 1} \right)^k \right]$$

- The translated and scaled polynomial is useful in the study of the conjugate gradient method:

$$T_k(x; a, b) = T_k\left(\frac{a + b - 2x}{b - a}\right)$$

where we have $|T_k(x; a, b)| \leq 1$ for all $x \in [a, b]$.



Convergence rate of Conjugate Gradient method

Theorem (Convergence rate of Conjugate Gradient method)

Let $\mathbf{A} \in \mathbb{R}^{n \times n}$ an SPD matrix then the **Conjugate Gradient** method converge to the solution $\mathbf{x}_* = \mathbf{A}^{-1}\mathbf{b}$ with at least linear r -rate in the norm $\|\cdot\|_{\mathbf{A}}$. Moreover we have the error estimate

$$\|\mathbf{e}_k\|_{\mathbf{A}} \lesssim 2 \left(\frac{\sqrt{\kappa} - 1}{\sqrt{\kappa} + 1} \right)^k \|\mathbf{e}_0\|_{\mathbf{A}}$$

$\kappa = M/m$ is the **condition number** where $m = \lambda_1$ is the smallest eigenvalue of \mathbf{A} and $M = \lambda_n$ is the biggest eigenvalue of \mathbf{A} .

The expression $a_k \lesssim b_k$ means that for all $\epsilon > 0$ there exists $k_0 > 0$ such that:

$$a_k \leq (1 - \epsilon)b_k, \quad \forall k > k_0$$



Proof.

From the estimate

$$\|\mathbf{e}_k\|_{\mathbf{A}} \leq \max_{\lambda \in [m, M]} |P(\lambda)| \|\mathbf{e}_0\|_{\mathbf{A}}, \quad P \in \mathbb{P}^k, P(0) = 1$$

choosing $P(x) = T_k(x; m, M)/T_k(0; m, M)$ from the fact that $|T_k(x; m, M)| \leq 1$ for $x \in [m, M]$ we have

$$\|\mathbf{e}_k\|_{\mathbf{A}} \leq T_k(0; m, M)^{-1} \|\mathbf{e}_0\|_{\mathbf{A}} = T_k \left(\frac{M+m}{M-m} \right)^{-1} \|\mathbf{e}_0\|_{\mathbf{A}}$$

observe that $\frac{M+m}{M-m} = \frac{\kappa+1}{\kappa-1}$ and

$$T_k \left(\frac{\kappa+1}{\kappa-1} \right)^{-1} = 2 \left[\left(\frac{\sqrt{\kappa}+1}{\sqrt{\kappa}-1} \right)^k + \left(\frac{\sqrt{\kappa}-1}{\sqrt{\kappa}+1} \right)^k \right]^{-1}$$

finally notice that $\left(\frac{\sqrt{\kappa}-1}{\sqrt{\kappa}+1} \right)^k \rightarrow 0$ as $k \rightarrow \infty$. □



Outline

- 1 The Steepest Descent iterative scheme
- 2 Conjugate direction method
- 3 Conjugate Gradient method
- 4 Conjugate Gradient convergence rate
- 5 **Preconditioning the Conjugate Gradient method**
- 6 Nonlinear Conjugate Gradient extension

Preconditioning

Problem (Preconditioned linear system)

Given $\mathbf{A}, \mathbf{P} \in \mathbb{R}^{n \times n}$, with \mathbf{A} an SPD matrix and \mathbf{P} non singular matrix and $\mathbf{b} \in \mathbb{R}^n$.

$$\text{Find } \mathbf{x}_* \in \mathbb{R}^n \text{ such that: } \mathbf{P}^{-T} \mathbf{A} \mathbf{x}_* = \mathbf{P}^{-T} \mathbf{b}.$$

A **good** choice for \mathbf{P} should be such that $\mathbf{M} = \mathbf{P}^T \mathbf{P} \approx \mathbf{A}$, where \approx denotes that \mathbf{M} is an approximation of \mathbf{A} in **some sense to precise later**.

Notice that:

- \mathbf{P} non singular imply:

$$\mathbf{P}^{-T} (\mathbf{b} - \mathbf{A} \mathbf{x}) = \mathbf{0} \iff \mathbf{b} - \mathbf{A} \mathbf{x} = \mathbf{0};$$

- \mathbf{A} SPD imply $\tilde{\mathbf{A}} = \mathbf{P}^{-T} \mathbf{A} \mathbf{P}^{-1}$ is also SPD (obvious proof).

Now we reformulate the preconditioned system:

Problem (Preconditioned linear system)

Given $\mathbf{A}, \mathbf{P} \in \mathbb{R}^{n \times n}$, with \mathbf{A} an SPD matrix and \mathbf{P} non singular matrix and $\mathbf{b} \in \mathbb{R}^n$ the preconditioned problem is the following:

$$\text{Find } \tilde{\mathbf{x}}_* \in \mathbb{R}^n \text{ such that: } \tilde{\mathbf{A}}\tilde{\mathbf{x}}_* = \tilde{\mathbf{b}}$$

where

$$\tilde{\mathbf{A}} = \mathbf{P}^{-T}\mathbf{A}\mathbf{P}^{-1} \quad \tilde{\mathbf{b}} = \mathbf{P}^{-T}\mathbf{b}$$

notice that if \mathbf{x}_* is the solution of the linear system $\mathbf{A}\mathbf{x} = \mathbf{b}$ then $\tilde{\mathbf{x}}_* = \mathbf{P}\mathbf{x}_*$ is the solution of the linear system $\tilde{\mathbf{A}}\tilde{\mathbf{x}} = \tilde{\mathbf{b}}$.



PCG: preliminary version

initial step:

$k \leftarrow 0$; \mathbf{x}_0 assigned;

$\tilde{\mathbf{x}}_0 \leftarrow \mathbf{P}\mathbf{x}_0$; $\tilde{\mathbf{r}}_0 \leftarrow \tilde{\mathbf{b}} - \tilde{\mathbf{A}}\tilde{\mathbf{x}}_0$; $\tilde{\mathbf{p}}_1 \leftarrow \tilde{\mathbf{r}}_0$;

while $\|\tilde{\mathbf{r}}_k\| > \epsilon$ **do**

$k \leftarrow k + 1$;

Conjugate direction method

$$\tilde{\alpha}_k \leftarrow \frac{\tilde{\mathbf{r}}_{k-1}^T \tilde{\mathbf{r}}_{k-1}}{\tilde{\mathbf{p}}_k^T \tilde{\mathbf{A}}\tilde{\mathbf{p}}_k};$$

$$\tilde{\mathbf{x}}_k \leftarrow \tilde{\mathbf{x}}_{k-1} + \tilde{\alpha}_k \tilde{\mathbf{p}}_k;$$

$$\tilde{\mathbf{r}}_k \leftarrow \tilde{\mathbf{r}}_{k-1} - \tilde{\alpha}_k \tilde{\mathbf{A}}\tilde{\mathbf{p}}_k;$$

Residual orthogonalization

$$\tilde{\beta}_k \leftarrow \frac{\tilde{\mathbf{r}}_k^T \tilde{\mathbf{r}}_k}{\tilde{\mathbf{r}}_{k-1}^T \tilde{\mathbf{r}}_{k-1}};$$

$$\tilde{\mathbf{p}}_{k+1} \leftarrow \tilde{\mathbf{r}}_k + \tilde{\beta}_k \tilde{\mathbf{p}}_k;$$

end while

final step

$$\mathbf{P}^{-1}\tilde{\mathbf{x}}_k;$$



Conjugate gradient algorithm applied to $\tilde{A}\tilde{x} = \tilde{b}$ require the evaluation of thing like:

$$\tilde{A}\tilde{p}_k = P^{-T}AP^{-1}\tilde{p}_k.$$

this can be done **without evaluate directly the matrix \tilde{A}** , by the following operations:

- ① solve $Ps'_k = \tilde{p}_k$ for $s'_k = P^{-1}\tilde{p}_k$;
- ② evaluate $s''_k = As'_k$;
- ③ solve $P^T s'''_k = s''_k$ for $s'''_k = P^{-T}s''_k$.

Step 1 and 3 require the solution of two auxiliary linear system. This is not a big problem if P and P^T are triangular matrices (see e.g. **incomplete Cholesky**).



However... we can reformulate the algorithm using only the matrices A and P !

Definition

For all $k \geq 1$, we introduce the vector $q_k = P^{-1}\tilde{p}_k$.

Observation

If the vectors $\tilde{p}_1, \tilde{p}_2, \dots, \tilde{p}_k$ for all $1 \leq k \leq n$ are \tilde{A} -conjugate, then the corresponding vectors q_1, q_2, \dots, q_k are A -conjugate.

In fact:

$$q_j^T A q_i = \underbrace{\tilde{p}_j^T}_{=q_j^T} P^{-T} A \underbrace{P^{-1}\tilde{p}_i}_{=q_i} = \tilde{p}_j^T \underbrace{\tilde{A}}_{=P^{-T}AP^{-1}} \tilde{p}_i = 0, \quad \text{if } i \neq j,$$

that is a consequence of \tilde{A} -conjugation of vectors \tilde{p}_i .



Definition

For all $k \geq 1$, we introduce the vectors

$$\mathbf{x}_k = \mathbf{x}_{k-1} + \tilde{\alpha}_k \mathbf{q}_k.$$

Observation

If we assume, *by construction*, $\tilde{\mathbf{x}}_0 = \mathbf{P}\mathbf{x}_0$, then we have

$$\tilde{\mathbf{x}}_k = \mathbf{P}\mathbf{x}_k, \quad \text{for all } k \text{ with } 1 \leq k \leq n.$$

In fact, if $\tilde{\mathbf{x}}_{k-1} = \mathbf{P}\mathbf{x}_{k-1}$ (inductive hypothesis), then

$$\begin{aligned} \tilde{\mathbf{x}}_k &= \tilde{\mathbf{x}}_{k-1} + \tilde{\alpha}_k \tilde{\mathbf{p}}_k && \text{[preconditioned CG]} \\ &= \mathbf{P}\mathbf{x}_{k-1} + \tilde{\alpha}_k \mathbf{P}\mathbf{q}_k && \text{[inductive Hyp. defs of } \mathbf{q}_k \text{]} \\ &= \mathbf{P}(\mathbf{x}_{k-1} + \tilde{\alpha}_k \mathbf{q}_k) && \text{[obvious]} \\ &= \mathbf{P}\mathbf{x}_k && \text{[defs. of } \mathbf{x}_k \text{]} \end{aligned}$$



Observation

Because $\tilde{\mathbf{x}}_k = \mathbf{P}\mathbf{x}_k$ for all $k \geq 0$, we have the recurrence between the corresponding residue $\tilde{\mathbf{r}}_k = \tilde{\mathbf{b}} - \tilde{\mathbf{A}}\tilde{\mathbf{x}}$ and $\mathbf{r}_k = \mathbf{b} - \mathbf{A}\mathbf{x}_k$:

$$\tilde{\mathbf{r}}_k = \mathbf{P}^{-T} \mathbf{r}_k.$$

In fact,

$$\begin{aligned} \tilde{\mathbf{r}}_k &= \tilde{\mathbf{b}} - \tilde{\mathbf{A}}\tilde{\mathbf{x}}_k, && \text{[defs. of } \tilde{\mathbf{r}}_k \text{]} \\ &= \mathbf{P}^{-T} \mathbf{b} - \mathbf{P}^{-T} \mathbf{A} \mathbf{P}^{-1} \mathbf{P}\mathbf{x}_k, && \text{[defs. of } \tilde{\mathbf{b}}, \tilde{\mathbf{A}}, \tilde{\mathbf{x}}_k \text{]} \\ &= \mathbf{P}^{-T} (\mathbf{b} - \mathbf{A}\mathbf{x}_k), && \text{[obvious]} \\ &= \mathbf{P}^{-T} \mathbf{r}_k. && \text{[defs. of } \mathbf{r}_k \text{]} \end{aligned}$$



Definition

For all k , with $1 \leq k \leq n$, the vector z_k is the solution of the linear system

$$Mz_k = r_k.$$

where $M = P^T P$. Formally,

$$z_k = M^{-1}r_k = P^{-1}P^{-T}r_k.$$

Using the vectors $\{z_k\}$,

- we can express $\tilde{\alpha}_k$ and $\tilde{\beta}_k$ in terms of A , the residual r_k , and conjugate direction q_k ;
- we can build a recurrence relation for the A -conjugate directions q_k .



Observation

$$\begin{aligned} \tilde{\alpha}_k &= \frac{\tilde{r}_{k-1}^T \tilde{r}_{k-1}}{\tilde{p}_k^T \tilde{A} \tilde{p}_k} = \frac{r_{k-1}^T P^{-1} P^{-T} r_{k-1}}{q_k^T P^T P^{-T} A P^{-1} P q_k} = \frac{r_{k-1}^T M^{-1} r_{k-1}}{q_k^T A q_k}, \\ &= \boxed{\frac{r_{k-1}^T z_{k-1}}{q_k^T A q_k}}. \end{aligned}$$

Observation

$$\begin{aligned} \tilde{\beta}_k &= \frac{\tilde{r}_k^T \tilde{r}_k}{\tilde{r}_{k-1}^T \tilde{r}_{k-1}} = \frac{r_k^T P^{-1} P^{-T} r_k}{r_{k-1}^T P^{-1} P^{-T} r_{k-1}} = \frac{r_k^T M^{-1} r_k}{r_{k-1}^T M^{-1} r_{k-1}}, \\ &= \boxed{\frac{r_k^T z_k}{r_{k-1}^T z_{k-1}}}. \end{aligned}$$



Observation

Using the vector $\mathbf{z}_k = \mathbf{M}^{-1}\mathbf{r}_k$, the following recurrence is true

$$\mathbf{q}_{k+1} = \mathbf{z}_k + \tilde{\beta}_k \mathbf{q}_k$$

In fact:

$$\tilde{\mathbf{p}}_{k+1} = \tilde{\mathbf{r}}_k + \tilde{\beta}_k \tilde{\mathbf{p}}_k \quad [\text{preconditioned CG}]$$

$$\mathbf{P}^{-1} \tilde{\mathbf{p}}_{k+1} = \mathbf{P}^{-1} \tilde{\mathbf{r}}_k + \tilde{\beta}_k \mathbf{P}^{-1} \tilde{\mathbf{p}}_k \quad [\text{left mult } \mathbf{P}^{-1}]$$

$$\mathbf{P}^{-1} \tilde{\mathbf{p}}_{k+1} = \mathbf{P}^{-1} \mathbf{P}^{-T} \mathbf{r}_k + \tilde{\beta}_k \mathbf{P}^{-1} \tilde{\mathbf{p}}_k \quad [\mathbf{r}_{k+1} = \mathbf{P}^{-T} \mathbf{r}_{k+1}]$$

$$\mathbf{P}^{-1} \tilde{\mathbf{p}}_{k+1} = \mathbf{M}^{-1} \mathbf{r}_k + \tilde{\beta}_k \mathbf{P}^{-1} \tilde{\mathbf{p}}_k \quad [\mathbf{M}^{-1} = \mathbf{P}^{-1} \mathbf{P}^{-T}]$$

$$\mathbf{q}_{k+1} = \mathbf{z}_k + \tilde{\beta}_k \mathbf{q}_k \quad [\mathbf{q}_k = \mathbf{P}^{-1} \tilde{\mathbf{p}}_k]$$



PCG: final version

initial step:

$k \leftarrow 0$; \mathbf{x}_0 assigned;

$\mathbf{r}_0 \leftarrow \mathbf{b} - \mathbf{A}\mathbf{x}_0$; $\mathbf{q}_1 \leftarrow \mathbf{r}_0$;

while $\|\mathbf{z}_k\| > \epsilon$ **do**

$k \leftarrow k + 1$;

Conjugate direction method

$$\tilde{\alpha}_k \leftarrow \frac{\mathbf{r}_{k-1}^T \mathbf{z}_{k-1}}{\mathbf{q}_k^T \mathbf{A} \mathbf{q}_k};$$

$$\mathbf{x}_k \leftarrow \mathbf{x}_{k-1} + \tilde{\alpha}_k \mathbf{q}_k;$$

$$\mathbf{r}_k \leftarrow \mathbf{r}_{k-1} - \tilde{\alpha}_k \mathbf{A} \mathbf{q}_k;$$

Preconditioning

$$\mathbf{z}_k = \mathbf{M}^{-1} \mathbf{r}_k;$$

Residual orthogonalization

$$\tilde{\beta}_k \leftarrow \frac{\mathbf{r}_k^T \mathbf{z}_k}{\mathbf{r}_{k-1}^T \mathbf{z}_{k-1}};$$

$$\mathbf{q}_{k+1} \leftarrow \mathbf{z}_k + \tilde{\beta}_k \mathbf{q}_k;$$

end while



Outline

- 1 The Steepest Descent iterative scheme
- 2 Conjugate direction method
- 3 Conjugate Gradient method
- 4 Conjugate Gradient convergence rate
- 5 Preconditioning the Conjugate Gradient method
- 6 Nonlinear Conjugate Gradient extension

Nonlinear Conjugate Gradient extension

- 1 The conjugate gradient algorithm can be extended for nonlinear minimization.
- 2 Fletcher and Reeves extend CG for the minimization of a general non linear function $f(\mathbf{x})$ as follows:
 - 1 Substitute the evaluation of α_k by an line search
 - 2 Substitute the residual \mathbf{r}_k with the gradient $\nabla f(\mathbf{x}_k)$
- 3 We also translate the index for the search direction \mathbf{p}_k to be more consistent with the gradients. The resulting algorithm is in the next slide

Fletcher and Reeves Nonlinear Conjugate Gradient

initial step:

$k \leftarrow 0$; \mathbf{x}_0 assigned;

$f_0 \leftarrow f(\mathbf{x}_0)$; $\mathbf{g}_0 \leftarrow \nabla f(\mathbf{x}_0)^T$;

$\mathbf{p}_0 \leftarrow -\mathbf{g}_0$;

while $\|\mathbf{g}_k\| > \epsilon$ **do**

$k \leftarrow k + 1$;

Conjugate direction method

Compute α_k by line-search;

$\mathbf{x}_k \leftarrow \mathbf{x}_{k-1} + \alpha_k \mathbf{p}_{k-1}$;

$\mathbf{g}_k \leftarrow \nabla f(\mathbf{x}_k)^T$;

Residual orthogonalization

$\beta_k^{FR} \leftarrow \frac{\mathbf{g}_k^T \mathbf{g}_k}{\mathbf{g}_{k-1}^T \mathbf{g}_{k-1}}$;

$\mathbf{p}_k \leftarrow -\mathbf{g}_k + \beta_k^{FR} \mathbf{p}_{k-1}$;

end while



- 1 To ensure convergence and apply Zoutendijk global convergence theorem we need to ensure that \mathbf{p}_k is a descent direction.
- 2 \mathbf{p}_0 is a descent direction by construction, for \mathbf{p}_k we have

$$\mathbf{g}_k^T \mathbf{p}_k = -\|\mathbf{g}_k\|^2 + \beta_k^{FR} \mathbf{g}_k^T \mathbf{p}_{k-1}$$

if the line-search is **exact** then $\mathbf{g}_k^T \mathbf{p}_{k-1} = 0$ because \mathbf{p}_{k-1} is the direction of the line-search. So by induction \mathbf{p}_k is a descent direction.

- 3 Exact line-search is expensive, however if we use inexact line-search with **strong Wolfe** conditions
 - 1 **sufficient decrease**: $f(\mathbf{x}_k + \alpha_k \mathbf{p}_k) \leq f(\mathbf{x}_k) + c_1 \alpha_k \nabla f(\mathbf{x}_k)^T \mathbf{p}_k$;
 - 2 **curvature condition**: $|\nabla f(\mathbf{x}_k + \alpha_k \mathbf{p}_k)^T \mathbf{p}_k| \leq c_2 |\nabla f(\mathbf{x}_k)^T \mathbf{p}_k|$.

with $0 < c_1 < c_2 < 1/2$ then we can prove that \mathbf{p}_k is a descent direction.




The previous consideration permits to say that Fletcher and Reeves nonlinear conjugate gradient method with strong Wolfe line-search is globally convergent¹

To prove globally convergence we need the following lemma:

Lemma (descent direction bound)

Suppose we apply Fletcher and Reeves nonlinear conjugate gradient method to $f(x)$ with strong Wolfe line-search with $0 < c_2 < 1/2$. The the method generates descent direction \mathbf{p}_k that satisfy the following inequality

$$-\frac{1}{1-c_2} \leq \frac{\mathbf{g}_k^T \mathbf{p}_k}{\|\mathbf{g}_k\|^2} \leq -\frac{1-2c_2}{1-c_2}, \quad k = 0, 1, 2, \dots$$

¹globally here means that Zoutendijk like theorem apply 

Proof. (1/3).

The proof is by induction. First notice that the function

$$t(\xi) = \frac{2\xi - 1}{1 - \xi}$$

is monotonically increasing on the interval $[0, 1/2]$ and that $t(0) = -1$ and $t(1/2) = 0$. Hence, because of $c_2 \in (0, 1/2)$ we have:

$$-1 < \frac{2c_2 - 1}{1 - c_2} < 0. \quad (\star)$$

base of induction $k = 0$: For $k = 0$ we have $\mathbf{p}_0 = -\mathbf{g}_0$ so that $\mathbf{g}_0^T \mathbf{p}_0 / \|\mathbf{g}_0\|^2 = -1$. From (\star) the lemma inequality is trivially satisfied.

Proof.

(2/3).

Using update direction formula's of the algorithm:

$$\beta_k^{FR} = \frac{\mathbf{g}_k^T \mathbf{g}_k}{\mathbf{g}_{k-1}^T \mathbf{g}_{k-1}} \quad \mathbf{p}_k = -\mathbf{g}_k + \beta_k^{FR} \mathbf{p}_{k-1}$$

we can write

$$\frac{\mathbf{g}_k^T \mathbf{p}_k}{\|\mathbf{g}_k\|^2} = -1 + \beta_k^{FR} \frac{\mathbf{g}_k^T \mathbf{p}_{k-1}}{\|\mathbf{g}_k\|^2} = -1 + \frac{\mathbf{g}_k^T \mathbf{p}_{k-1}}{\|\mathbf{g}_{k-1}\|^2}$$

and by using second strong Wolfe condition:

$$-1 + c_2 \frac{\mathbf{g}_{k-1}^T \mathbf{p}_{k-1}}{\|\mathbf{g}_{k-1}\|^2} \leq \frac{\mathbf{g}_k^T \mathbf{p}_k}{\|\mathbf{g}_k\|^2} \leq -1 - c_2 \frac{\mathbf{g}_{k-1}^T \mathbf{p}_{k-1}}{\|\mathbf{g}_{k-1}\|^2}$$



Proof.

(3/3).

by induction we have

$$\frac{1}{1 - c_2} \geq -\frac{\mathbf{g}_{k-1}^T \mathbf{p}_{k-1}}{\|\mathbf{g}_{k-1}\|^2} > 0$$

so that

$$\frac{\mathbf{g}_k^T \mathbf{p}_k}{\|\mathbf{g}_k\|^2} \leq -1 - c_2 \frac{\mathbf{g}_{k-1}^T \mathbf{p}_{k-1}}{\|\mathbf{g}_{k-1}\|^2} \leq -1 + c_2 \frac{1}{1 - c_2} = \frac{2c_2 - 1}{1 - c_2}$$

and

$$\frac{\mathbf{g}_k^T \mathbf{p}_k}{\|\mathbf{g}_k\|^2} \geq -1 + c_2 \frac{\mathbf{g}_{k-1}^T \mathbf{p}_{k-1}}{\|\mathbf{g}_{k-1}\|^2} \geq -1 - c_2 \frac{1}{1 - c_2} = -\frac{1}{1 - c_2}$$

□



- 1 The inequality of the the previous lemma can be written as:

$$\frac{1}{1 - c_2} \frac{\|\mathbf{g}_k\|}{\|\mathbf{p}_k\|} \geq -\frac{\mathbf{g}_k^T \mathbf{p}_k}{\|\mathbf{g}_k\| \|\mathbf{p}_k\|} \geq \frac{1 - 2c_2}{1 - c_2} \frac{\|\mathbf{g}_k\|}{\|\mathbf{p}_k\|} > 0$$

- 2 Remembering the Zoutendijk theorem we have

$$\sum_{k=1}^{\infty} (\cos \theta_k)^2 \|\mathbf{g}_k\|^2 < \infty, \quad \text{where} \quad \cos \theta_k = -\frac{\mathbf{g}_k^T \mathbf{p}_k}{\|\mathbf{g}_k\| \|\mathbf{p}_k\|}$$

- 3 so that if $\|\mathbf{g}_k\| / \|\mathbf{p}_k\|$ is bounded from below we have that $\cos \theta_k \geq \delta$ for all k and then from Zoutendijk theorem the scheme converge.
- 4 Unfortunately this bound cant be proved so that Zoutendijk theorem cant be applied directly. However it is possible to prove a weaker results, i.e. that $\liminf_{k \rightarrow \infty} \|\mathbf{g}_k\| = 0!$



Convergence of Fletcher and Reeves method

Assumption (Regularity assumption)

We assume $f \in C^1(\mathbb{R}^n)$ with Lipschitz continuous gradient, i.e. there exists $\gamma > 0$ such that

$$\|\nabla f(\mathbf{x})^T - \nabla f(\mathbf{y})^T\| \leq \gamma \|\mathbf{x} - \mathbf{y}\|, \quad \forall \mathbf{x}, \mathbf{y} \in \mathbb{R}^n$$



Theorem (Convergence of Fletcher and Reeves method)

Suppose the method of **Fletcher and Reeves** is implemented with strong Wolfe line-search with $0 < c_1 < c_2 < 1/2$. If $f(x)$ and x_0 satisfy the previous regularity assumptions, then

$$\liminf_{k \rightarrow \infty} \|g_k\| = 0$$

Proof.

(1/4).

From previous Lemma we have

$$\cos \theta_k \geq \frac{1}{1 - c_2} \frac{\|g_k\|}{\|p_k\|} \quad k = 1, 2, \dots$$

substituting in Zoutendijk condition we have $\sum_{k=1}^{\infty} \frac{\|g_k\|^4}{\|p_k\|^2} < \infty$.

The proof is by contradiction. in fact if theorem is not true than the series diverge. Next we want to bound $\|p_k\|$.



Proof. (bounding $\|p_k\|$)

(2/4).

Using second Wolfe condition and previous Lemma

$$|g_k^T p_{k-1}| \leq -c_2 g_k^T p_{k-1} \leq \frac{c_2}{1 - c_2} \|g_{k-1}\|^2$$

using $p_k = -g_k + \beta_k^{FR} p_{k-1}$ we have

$$\begin{aligned} \|p_k\|^2 &\leq \|g_k\|^2 + 2\beta_k^{FR} |g_k^T p_{k-1}| + (\beta_k^{FR})^2 \|p_{k-1}\|^2 \\ &\leq \|g_k\|^2 + \frac{2c_2}{1 - c_2} \beta_k^{FR} \|g_{k-1}\|^2 + (\beta_k^{FR})^2 \|p_{k-1}\|^2 \end{aligned}$$

recall that $\beta_k^{FR} = \|g_k\|^2 / \|g_{k-1}\|^2$ then

$$\|p_k\|^2 \leq \frac{1 + c_2}{1 - c_2} \|g_k\|^2 + (\beta_k^{FR})^2 \|p_{k-1}\|^2$$



Proof. (bounding $\|\mathbf{p}_k\|$) (3/4).

setting $c_3 = \frac{1+c_2}{1-c_2}$ and using repeatedly the last inequality we obtain:

$$\begin{aligned}
 \|\mathbf{p}_k\|^2 &\leq c_3 \|\mathbf{g}_k\|^2 + (\beta_k^{FR})^2 (c_3 \|\mathbf{g}_{k-1}\|^2 + (\beta_{k-1}^{FR})^2 \|\mathbf{p}_{k-2}\|^2) \\
 &= c_3 \|\mathbf{g}_k\|^4 \left(\|\mathbf{g}_k\|^{-2} + \|\mathbf{g}_{k-1}\|^{-2} \right) + \frac{\|\mathbf{g}_k\|^4}{\|\mathbf{g}_{k-2}\|^4} \|\mathbf{p}_{k-2}\|^2 \\
 &\leq c_3 \|\mathbf{g}_k\|^4 \left(\|\mathbf{g}_k\|^{-2} + \|\mathbf{g}_{k-1}\|^{-2} + \|\mathbf{g}_{k-2}\|^{-2} \right) \\
 &\quad + \frac{\|\mathbf{g}_k\|^4}{\|\mathbf{g}_{k-3}\|^4} \|\mathbf{p}_{k-3}\|^2 \\
 &\leq c_3 \|\mathbf{g}_k\|^4 \sum_{j=1}^k \|\mathbf{g}_j\|^{-2}
 \end{aligned}$$



Proof. (4/4).

Suppose now **by contradiction** there exists $\delta > 0$ such that $\|\mathbf{g}_k\| \geq \delta$ ^a by using the regularity assumptions we have

$$\|\mathbf{p}_k\|^2 \leq c_3 \|\mathbf{g}_k\|^4 \sum_{j=1}^k \|\mathbf{g}_j\|^{-2} \leq c_3 \|\mathbf{g}_k\|^4 \delta^{-2} k$$

Substituting in Zoutendijk condition we have

$$\infty > \sum_{k=1}^{\infty} \frac{\|\mathbf{g}_k\|^4}{\|\mathbf{p}_k\|^2} \geq \frac{\delta^2}{c_4} \sum_{k=1}^{\infty} \frac{1}{k} = \infty$$

this contradict assumption. □

^athe correct assumption is that there exists k_0 such that $\|\mathbf{g}_k\| \geq \delta$ for $k \geq k_0$ but this complicate a little bit the following inequality without introducing new idea.



Weakness of Fletcher and Reeves method

- Suppose that \mathbf{p}_k is a **bad** search direction, i.e. $\cos \theta_k \approx 0$.
- From the **descent direction bound** Lemma (see slide 91) we have

$$\frac{1}{1 - c_2} \frac{\|\mathbf{g}_k\|}{\|\mathbf{p}_k\|} \geq \cos \theta_k \geq \frac{1 - 2c_2}{1 - c_2} \frac{\|\mathbf{g}_k\|}{\|\mathbf{p}_k\|} > 0$$

- so that to have $\cos \theta_k \approx 0$ we need $\|\mathbf{p}_k\| \gg \|\mathbf{g}_k\|$.
- since \mathbf{p}_k is a bad direction near orthogonal to \mathbf{g}_k it is likely that the step is small and $\mathbf{x}_{k+1} \approx \mathbf{x}_k$. If so we have also $\mathbf{g}_{k+1} \approx \mathbf{g}_k$ and $\beta_{k+1}^{FR} \approx 1$.
- but remember that $\mathbf{p}_{k+1} \leftarrow -\mathbf{g}_{k+1} + \beta_{k+1}^{FR} \mathbf{p}_k$, so that $\mathbf{p}_{k+1} \approx \mathbf{p}_k$.
- This means that a **long sequence of unproductive iterates** will follow.



Polack and Ribière Nonlinear Conjugate Gradient

- 1 The previous problem can be elided if we restart anew when the iterate stagnate.
- 2 Restarting is obtained by simply set $\beta_k^{FR} = 0$.
- 3 A more elegant solution can be obtained with a new definition of β_k due to Polack and Ribière is the following:

$$\beta_k^{PR} = \frac{\mathbf{g}_k^T (\mathbf{g}_k - \mathbf{g}_{k-1})}{\mathbf{g}_{k-1}^T \mathbf{g}_{k-1}}$$

- 4 This definition of β_k^{PR} is identical of β_k^{FR} in the case of quadratic function because $\mathbf{g}_k^T \mathbf{g}_{k-1} = 0$. The definition **differs** in non linear case and in particular when there is stagnation i.e. $\mathbf{g}_k \approx \mathbf{g}_{k-1}$ we have $\beta_k^{PR} \approx 0$, i.e. we have an **automatic restart**.



Polack and Ribière Nonlinear Conjugate Gradient

initial step:

$k \leftarrow 0$; \mathbf{x}_0 assigned;

$f_0 \leftarrow f(\mathbf{x}_0)$; $\mathbf{g}_0 \leftarrow \nabla f(\mathbf{x}_0)^T$;

$\mathbf{p}_0 \leftarrow -\mathbf{g}_0$;

while $\|\mathbf{g}_k\| > \epsilon$ **do**

$k \leftarrow k + 1$;

Conjugate direction method

Compute α_k by line-search;

$\mathbf{x}_k \leftarrow \mathbf{x}_{k-1} + \alpha_k \mathbf{p}_{k-1}$;

$\mathbf{g}_k \leftarrow \nabla f(\mathbf{x}_k)^T$;

Residual orthogonalization

$$\beta_k^{PR} \leftarrow \frac{\mathbf{g}_k^T (\mathbf{g}_k - \mathbf{g}_{k-1})}{\mathbf{g}_{k-1}^T \mathbf{g}_{k-1}};$$

$$\mathbf{p}_k \leftarrow -\mathbf{g}_k + \beta_k^{PR} \mathbf{p}_{k-1};$$

end while



Weakness of Polack and Ribière method

(1/2)

- Although the modification is minimal, for the Polack and Ribière method with strong Wolfe line-search it can happen that \mathbf{p}_k is not a descent direction.
- If \mathbf{p}_k is not a descent direction we can restart i.e. set $\beta_k^{PR} = 0$ or modify β_k^{PR} as follows

$$\beta_k^{PR+} = \max\{\beta_k^{PR}, 0\}$$

this new coefficient with a modified Wolfe line-search ensure that \mathbf{p}_k is a descent direction.



Weakness of Polack and Ribière method

(2/2)

- Polack and Ribière choice on the average perform better than Fletcher and Reeves but there is **not** convergence results!
- Although there is not convergence results there is a negative results due to Powell:

Theorem

Consider the Polack and Ribière method with exact line-search. There exists a twice continuously differentiable function $f : \mathbb{R}^3 \mapsto \mathbb{R}$ and a starting point x_0 such that the sequence of gradients $\{ \|g_k\| \}$ is bounded away from zero.

- However in spite of this results Polack and Ribière is the first choice among conjugate direction methods.



Other choices




- There are many other modification of the coefficient β_k that collapse to the same coefficient in the case of quadratic function. One important choice is the Hestenes and Stiefel choice

$$\beta_k^{HS} = \frac{\mathbf{g}_k^T (\mathbf{g}_k - \mathbf{g}_{k-1})}{(\mathbf{g}_k^T - \mathbf{g}_{k-1}^T) \mathbf{p}_{k-1}}$$

- For this choice there is similar convergence results of Fletcher and Reeves and similar performance.



References

-  J. E. Dennis, Jr. and Robert B. Schnabel
Numerical Methods for Unconstrained Optimization and
Nonlinear Equations
SIAM, Classics in Applied Mathematics, **16**, 1996.
-  J. Nocedal and S. J. Wright
Numerical Optimization
Springer Series in Operation Research, 1999.
-  J. Stoer and R. Bulirsch
Introduction to numerical analysis
Springer-Verlag, Texts in Applied Mathematics, **12**, 2002.